

One of the arguments against the findings of conventional population studies is the fact that smokers are self-selected and that other factors may be involved in the causation of IHD.

It has been maintained by Friedman & Rosenman (1972) that persons with a particular behavior pattern, type A, characterized by aggressiveness, striving and time consciousness, more often develop myocardial infarction than the more placid type B; and the type A person is also more likely to be a heavy smoker.

Questionnaire studies on the Swedish as well as the U.S. Twin Registries disclosed an association for males between smoking and angina pectoris when the results were analysed on a non-pair basis, but comparisons within smoking discordant pairs showed no association with smoking for the MZ twins (report from symposium in San Juan, Puerto Rico, 1969). These results are supported by a clinical study on 196 smoking discordant pairs (Lundman, 1966) from the Swedish Twin Registry, no difference being found with regard to IHD within the smoking discordant pairs.

Material: see Chapter I.

#### Methods

Information about smoking habits was obtained from questionnaire replies received in 1967/1970. If the twins had answered in 1967 as well as 1970, the latter report was used.

#### Results

Smoking habits in the male and female death discordant pairs according to the questionnaires are shown in Tables 26-28. When present and former smokers are combined (Table 28), the percentage distribution of male smokers is somewhat, but not significantly, higher for the IHD compared to the not-IHD death discordant pairs. Of the male twins who died from IHD, 83 % were smokers as against 85 % of their surviving co-twins, while 76 % of those who died from other causes than IHD were smokers compared with 72 % of their surviving co-twins. The corresponding figures for cigarette smokers in the IHD death discordant pairs are 50 % and 55 % compared to 40 % and 52 % among the not-IHD death discordant pairs. Nor are these differences significant.

There were more former cigarette smokers (Table 26) among the deceased male twins than the surviving co-twins, especially among those who died from IHD.

With respect to zygosity groups (Table 28), 60 % (6 of 10) of the

Table 26. Smoking habits in male death discordant pairs according to earlier questionnaires. Percentage distribution.

IHD		not IHD		Tot.		Tot.	
MZ	DZ	MZ	DZ	MZ	DZ	MZ	DZ
S	S	S	S	S	S	S	S
D	D	D	D	D	D	D	D
Tot.		Tot.		Tot.		Tot.	

Non smokers

Table 26. Smoking habits in male death discordant pairs according to earlier questionnaires. Percentage distribution.

	IHD								not IHD							
	MZ		DZ		XZ		Tot.		MZ		DZ		XZ		Tot.	
	D	S	D	S	D	S	D	S	D	S	D	S	D	S	D	S
Non smokers	10	20	20	16	20	0	18	15	20	16	26	19	0	0	24	18
Total present smokers	40	80	52	68	60	60	50	70	56	60	52	67	100	100	54	65
Total cigarette smokers	30	80	24	24	60	60	30	43	12	32	26	36	100	100	22	35
1-10 cigarettes daily	20	30	16	12	40	20	20	18	8	28	14	21	100	0	13	24
>11 cigarettes daily	10	50	8	12	20	40	10	25	4	4	12	14	0	100	9	12
Cigar and/or pipe	10	0	28	44	0	0	20	28	44	28	26	31	0	0	32	29
Total former smokers	50	0	28	16	20	40	33	15	24	24	21	14	0	0	22	18
Total cigarette smokers	30	0	16	12	20	40	20	13	24	20	14	14	0	0	18	16
1-10 cigarettes daily	10	0	16	8	20	20	15	8	16	16	10	14	0	0	12	15
>11 cigarettes daily	20	0	0	4	0	20	5	5	8	4	5	0	0	0	6	1
Cigar and/or pipe	20	0	12	4	0	0	13	3	0	4	7	0	0	0	4	1
Total number	10	10	25	25	5	5	40	40	25	25	42	42	1	1	68	68

D=Deceased twin; S=Surviving co-twin

Table 27. Smoking habits in female death discordant pairs according to earlier questionnaires. Percentage distribution.

	IHD								not IHD							
	MZ		DZ		XZ		Tot.		MZ		DZ		XZ		Tot.	
	D	S	D	S	D	S	D	S	D	S	D	S	D	S	D	S
Non smokers	75	50	78	67	-	-	76	59	70	77	85	79	50	100	79	79
Total present smokers	25	13	11	33	-	-	18	24	30	23	15	17	50	0	21	19
Total cigarette smokers	25	13	11	33	-	-	18	24	27	23	15	17	50	0	20	19
1-10 cigarettes daily	13	13	11	11	-	-	12	12	20	23	10	10	50	0	15	15
≥11 cigarettes daily	13	0	0	22	-	-	6	12	7	0	4	6	0	0	5	4
Cigar and/or pipe	0	0	0	0	-	-	0	0	3	0	0	0	0	0	1	0
Total former smokers	0	38	11	0	-	-	6	18	0	0	0	4	0	0	0	3
Total cigarette smokers	0	38	11	0	-	-	6	18	0	0	0	2	0	0	0	1
1-10 cigarettes daily	0	25	11	0	-	-	6	12	0	0	0	2	0	0	0	1
≥11 cigarettes daily	0	13	0	0	-	-	0	6	0	0	0	0	0	0	0	0
Cigar and/or pipe	0	0	0	0	-	-	0	0	0	0	0	2	0	0	0	1
Total number	8	8	9	9	0	0	17	17	30	30	48	48	2	2	80	80

D=Deceased twin; S=Surviving co-twin

Table 28. Smoking habits in male and female death discordant pairs according to earlier questionnaires with present and former smokers combined. Percentage distribution.

	IHD								not IHD							
	MZ		DZ		XZ		Tot.		MZ		DZ		XZ		Tot.	
	D	S	D	S	D	S	D	S	D	S	D	S	D	S	D	S

2028559362

Table 28. Smoking habits in male and female death discordant pairs according to earlier questionnaires with present and former smokers combined. Percentage distribution.

	IHD								not IHD							
	MZ		DZ		XZ		Tot.		MZ		DZ		XZ		Tot.	
	D	S	D	S	D	S	D	S	D	S	D	S	D	S	D	S
<b>MALES:</b>																
Non smokers	10	20	20	16	20	0	18	15	20	16	26	19	0	0	24	18
Total smokers	90	80	80	84	80	100	83	85	80	84	74	81	100	100	76	72
Total cigarette smokers	60	80	40	36	80	100	50	55	36	52	41	50	100	100	40	52
1-10 cigarettes daily	30	30	32	20	60	40	35	25	24	44	24	36	100	0	25	38
≥11 cigarettes daily	30	50	8	16	20	60	15	30	12	8	17	14	0	100	15	13
Cigar and/or pipe	30	0	40	48	0	0	33	30	44	32	33	31	0	0	37	31
Total number	10	10	25	25	5	5	40	40	25	25	42	42	1	1	68	68
<b>FEMALES:</b>																
Non smokers	75	50	78	67	-	-	77	59	70	77	85	79	50	100	79	79
Total smokers	25	50	22	33	-	-	24	41	30	23	15	21	50	0	21	21
Total cigarette smokers	25	50	22	33	-	-	24	41	30	23	15	19	50	0	20	20
1-10 cigarettes daily	13	38	22	11	-	-	18	35	30	23	15	19	50	0	15	20
≥11 cigarettes daily	13	13	0	22	-	-	6	18	7	0	4	6	0	0	3	4
Cigar and/or pipe	0	0	0	0	-	-	0	0	3	0	0	2	0	0	1	1
Total number	8	8	9	9	0	0	17	17	30	30	48	48	2	2	80	80

D=Deceased twin; S=Surviving co-twin

Table 29. Distribution of death discordant male twin pairs according to concordance/discordance with respect to smoking habits (present and former smokers combined).

		Deceased twin															
		IHD								not IHD							
		MZ		DZ		XZ		Tot.		MZ		DZ		XZ		Tot.	
		NS	SM	NS	SM	NS	SM	NS	SM	NS	SM	NS	SM	NS	SM	NS	SM
		NS	SM	NS	SM	NS	SM	NS	SM	NS	SM	NS	SM	NS	SM	NS	SM
Surviving co-twin	NS	1	1	2	2	0	0	3	3	4	0	4	4	0	0	8	4
	SM	0	8	3	18	1	4	4	30	1	20	7	27	0	1	8	48

Table 30. Distribution of death discordant male twin pairs according to concordance/discordance with respect to cigarette habits (present and former smokers combined).

		Deceased twin															
		IHD								not IHD							
		MZ		DZ		XZ		Tot.		MZ		DZ		XZ		Tot.	
		NS	SM	NS	SM	NS	SM	NS	SM	NS	SM	NS	SM	NS	SM	NS	SM
Surviving co-twin	NS	1	1	2	1	0	0	3	2	4	0	4	3	0	0	8	3
	SM	0	8	1	12	1	4	2	24	1	14	5	20	0	1	6	35

Table 31. Distribution of death discordant female twin pairs according to concordance/discordance with respect to smoking habits (present and former smokers combined).

		Deceased twin															
		IHD								not IHD							
		MZ		DZ		XZ		Tot.		MZ		DZ		XZ		Tot.	
		NS	SM	NS	SM	NS	SM	NS	SM	NS	SM	NS	SM	NS	SM	NS	SM
		NS	SM	NS	SM	NS	SM	NS	SM	NS	SM	NS	SM	NS	SM	NS	SM
Surviving co-twin	NS	4	0	6	0	0	0	10	0	19	4	35	3	1	1	55	8
	SM	2	2	1	2	0	0	3	4	2	5	6	4	0	0	8	9

SM=Smoker; NS=Non Smoker

Comments

MZ twins who died their surviving counterparts the corresponding daily is significant MZ co-twins, with pairs whose parents were smokers (50% vs 8%). However, with male MZ co-twin pairs, there is a surviving co-twin compared to the deceased partner. Of the few compared with figures for the twins. Nearly all the smokers, only differences were noted. Smoking did not cause death in Tables 29-30 of the smoking compared to 41 cigarette smokers. Figures for the twins there statistically respect to smoking.

MZ twins who died from IHD were cigarette smokers as against 80 % of their surviving co-twins. Counting only present smokers (Table 26), the corresponding figures are 30 % vs 80 %. Smoking >10 cigarettes daily is significantly ( $p < 0.05$ ) more common among the surviving male MZ co-twins, whose partners died from IHD, than among their counterparts whose partners died from other causes; this is true for present smokers (50 % vs 4 %) and for present and former smokers combined (50 % vs 8 %). The corresponding comparisons between the deceased male MZ co-twins give the same trend but it is not so pronounced. However, with regard to the larger group of death discordant male DZ pairs, there is a higher proportion of cigarette smokers among the surviving co-twins whose partners died from other causes (50 %) compared to those whose partners died from IHD (36 %), but among their deceased partners the proportions are about the same (41 % vs 40 %).

Of the female twins who died from IHD (Table 28), 24 % were smokers compared with 41 % of their surviving co-twins, while the corresponding figures for the not-IHD death discordant twins are 21 % and 21 %. Nearly all the female smokers were cigarette smokers. Of the former smokers, only one was found among the deceased co-twins. None of the differences recorded among the females is statistically significant.

Smoking discordance among the male death discordant pairs is shown in Tables 29-30. Only among the DZ twin pairs in whom death discordance was not caused by IHD is there an apparent smoking discordance: in 7 of the smoking discordant pairs the surviving co-twins were smokers compared to 4 pairs in which the deceased twins were smokers. If only cigarette smoking discordant pairs are considered, the corresponding figures for the not-IHD death discordant pairs were 5 and 3. Nor were there statistically significant differences among the females with respect to smoking discordance (Table 31).

#### Comments

With regard to mortality from IHD, especially in men, most studies indicate a positive relation to smoking (Doll & Hill, 1964; Best, 1966; Hammond, 1966; Kannel, 1966; Fletcher & Horn, 1970). The present number of smoking discordant twin pairs among the IHD death discordant pairs is too small to permit an evaluation of the effect of smoking on IHD mortality, but it can be mentioned that in the mortality study on the whole of the Swedish Twin Registry (Friberg et al, 1973), a hypermortality, partly due to IHD, was found among DZ smoking discordant pairs but not among MZ pairs in both men and women. It may well be that cigarette smoking only reflects a certain constitution and that

The findings of proportionally more cigarette smokers among the surviving male MZ co-twins whose partners died from IHD compared to those whose partners died from other causes could indicate an association between cigarette smoking and IHD. The larger male DZ group did not show similar differences in cigarette smoking habits. However, the deceased male MZ twins also displayed the same trend in cigarette smoking habits as the surviving co-twins, which may indicate that it is constitutional factors which lie behind these findings.

Studies on the association between alcohol consumption and IHD have yielded conflicting results. Chronic alcoholics have been considered to possess protection from coronary atherosclerosis. Several authors (e.g. Grant et al, 1959; Stare, 1961; Björck, 1963) found no association between alcohol consumption and IHD. Data from the Framingham study support the opinion that alcohol is not correlated with IHD (Kannel, 1966) - alcohol consumption apparently showed no association with the development of IHD, neither did it protect against atherosclerosis of the heart.

Material: see Chapter I.

[illegible]

Table 32. Distribution of death discordant male twin pairs according to registration in Alcohol Registry.

	IHD								not IHD							
	MZ		DZ		XZ		Tot.		MZ		DZ		XZ		Tot.	
	D	S	D	S	D	S	D	S	D	S	D	S	D	S	D	S
Registered	1	0	5	3	1	2	7	5	6	8	19	15	0	0	25	23
Total number	10	10	25	25	5	5	40	40	25	25	42	42	1	1	68	68

D=Deceased twin; S=Surviving co-twin

Table 33. Distribution of death discordant male twin pairs according to concordance/discordance with respect to registration in Alcohol Registry.

	Deceased twin																
	IHD								not IHD								
	MZ		DZ		XZ		Tot.		MZ		DZ		XZ		Tot.		
	NR	R	NR	R	NR	R	NR	R	NR	R	NR	R	NR	R	NR	R	
Surviving co-twin	NR	9	1	18	4	2	1	29	6	15	2	16	11	1	0	32	13
	R	0	0	2	1	2	0	4	1	4	4	7	8	0	0	11	12

R=Registered; NR=Not Registered



Persons with alcohol problems were defined operationally as those entered in a nationwide registry for misconduct while under the influence of alcoholic beverages. This registry has existed since 1932. All the male death discordant twins were matched with the registry.

The numbers of death discordant male twins with entries in the Alcohol Registry are shown in Table 32. Entries were found for 32 out of 108 (30 %) deceased twins compared to 28 out of 108 (26 %) surviving co-twins. Of the 40 twins who had died from IHD, 7 (18 %) had been registered compared with 5 of their surviving partners (13 %), whereas of the 68 who had died from other causes than IHD, 25 (37 %) had been registered compared with 23 of their surviving partners (34 %). A distribution of the registered twins in the death discordant pairs (Table 33) shows that only one of the IHD death discordant pairs, a DZ pair, is concordant with respect to registration compared to 12 in the not-IHD group. Among the discordant (registered/not registered) pairs there are no significant differences within the death discordant pairs.

An entry in the Alcohol Registry usually indicates that the person in question has alcohol problems. It was found by Helander (1972), that 11.5 % of male twins born 1900-1924 had at least one conviction for drunkenness compared to only 0.2 % of the female twins, the corresponding figure for male twins born 1925-1954 being 11.3 % and for females 0.6 %. Because so few female twins have been registered, only male twins were matched with the Alcohol Registry. A mortality follow-up of 9,000 twin pairs from the Swedish Twin Registry has shown entry in the Alcohol Registry to be associated with a higher mortality regardless of smoking and it was also found that 10 % of non-smokers were registered as against 30 % of those who smoked more than 10 cigarettes a day (Friberg et al, 1973). Concerning the cause-specific death rates, a weak association was found between registration and IHD. In the present investigation too, registration was somewhat more common among the deceased twins irrespective of the cause of death (IHD/not IHD). However, alcohol registration does not seem to have contributed to the high prevalence of IHD manifestations in the surviving co-twins in the IHD death discordant pairs. Using questionnaire replies in 1967 concerning

Various  
have been cons  
complexity of  
their investig  
evaluate. Stam  
urbanized soci  
Results dis  
economic groups  
lence of myocar

alcohol consumption, Myrhed (1974) investigated 70 male twin pairs discordant with respect to alcohol consumption. The alcohol discordant pairs did not differ in IHD manifestations evaluated as ST depressions in connection with exercise. A check of the questionnaire replies from 1967/1970 on alcohol consumption for the present material yielded a fairly high proportion of non respondents (27 % among the deceased male twins and 36 % among their surviving co-twins, compared to 44 % among the deceased female twins and 46 % among their surviving co-twins), which is why this source of information on alcohol consumption could not be used here.

#### PHYSICAL INACTIVITY AND SOME SOCIAL FACTORS

Physical inactivity or sedentary living has also been implicated as a risk factor for IHD (Simborg, 1970; Karvonen, 1972; Stamler, 1973). The findings are not as clearcut or consistent as for many other risk factors, probably due to the difficulty of defining. Results from the Health Insurance Plan (Frank et al, 1966) and the Framingham study (Kannel, 1966) have shown a striking increase of fatal myocardial infarction among physically inactive men. No such correlation was found to angina pectoris and coronary insufficiency in the Framingham study but the Health Insurance Plan showed angina pectoris to be even more common in the more active group of men. The Western collaborative study (Frank et al, 1966) has also confirmed that regular exercise habits were less common among those with a fatal as opposed to a non-fatal myocardial infarction. On the other hand, the seven countries study (Keys et al, 1970) disclosed no association between the incidence of IHD and physical inactivity in any of the cohorts. It is also worth noting that the members of the cohorts from Finland had the highest incidence of IHD but were usually the most physically active and non-obese.

Various social and psychosocial factors have been considered as risks for IHD (cf. Jenkins, 1971). The complexity of these variables and the lack of uniform methods for their investigation make such studies difficult to perform and evaluate. Stamler (1973) considers the modern way of life in highly urbanized societies to play an important role in the causation of IHD.

Results differ with regard to the correlation of different socioeconomic groups to IHD (Hofvendahl & Helmers, 1973). A higher prevalence of myocardial infarction has been noted among employers,

especially owners of small shops and salesmen (Börck et al, 1958). Analyses of later studies have tended instead towards a higher prevalence and incidence of IHD in the lower socio-economic groups (Antonowsky, 1968).

A low educational background has also been associated with an increased risk of myocardial infarction (Antonowsky, 1968; Hinkle, 1969). Social group mobility (Wardwell, 1968) as well as occupational mobility (Syme, 1964) are other sociologic factors which have been found to discriminate IHD patients from other subjects.

According to Friedman & Rosenman (1959), individuals exhibiting an emotional complex characterized by an excessive sense of time urgency, drive, and competitiveness (type A behavior) are more prone to the onset of IHD than are individuals of the converse type of behavior (type B). Type A subjects usually have elevated cholesterol values and are also more often heavy smokers than type B subjects. Other psychosocial characteristics found in men with myocardial infarction seem to be job dissatisfaction (Sales & House, 1971; Theorell, 1971), and overtime work (Kasanen et al, 1963; Liljefors, 1970).

Material: see Chapter I.

#### Methods

Information on physical inactivity and some social factors (change of place of work, extra work and place of residence) was obtained from questionnaires mailed to all twins in 1967 and 1970, while information on education was obtained from questionnaires in 1962. The twins were asked among other things if they had changed their place of work after the age of 25 and how often, if they had had extra work beside their ordinary job and if this happened  $\geq$  periodically. They were also asked about their place of residence after the age of 25 and if this had been mostly in big cities. Higher education is taken here to be education above the compulsory school, including folk-high school, vocational school and evening courses. The twins were also asked about how much exercise they had taken when 25-50 years of age; the answer "hardly any" was recorded here.

#### Results

The distribution of death discordant pairs according to their answers on physical inactivity and various sociologic variables is given in Tables 34 and 35. Hardly any exercise when 25-50 years of age is

Table 34. Physical inactivity and some sociologic variables in the male death discordant pairs according to earlier questionnaires. Absolute numbers and prevalence rates in % (parenthesis).

	IHD				not IHD				Tot.			
	MZ	DZ	XZ	Tot.	MZ	DZ	XZ	Tot.	MZ	DZ	XZ	Tot.
Hardly any exercise	2	1	0	3	6	0	0	6	3	2	4	9
	(100)	(100)	(0)	(33)	(100)	(0)	(0)	(100)	(33)	(67)	(67)	(78)

Table 34. Physical inactivity and some sociologic variables in the male death discordant pairs according to earlier questionnaires. Absolute numbers and prevalence rates in % (parenthesis).

	IHD								not IHD							
	MZ		DZ		XZ		Tot.		MZ		DZ		XZ		Tot.	
	D	S	D	S	D	S	D	S	D	S	D	S	D	S	D	S
Hardly any exercise	2 (20)	1 (10)	6 (24)	6 (24)	0 (0)	0 (0)	8 (20)	7 (18)	3 (12)	2 (8)	4 (10)	5 (12)	0 (0)	0 (0)	7 (10)	7 (10)
Change of place of work $\geq$ 5 times	0 (0)	0 (0)	0 (0)	2 (8)	3 (60)	1 (20)	3 (8)	3 (18)	4 (16)	4 (16)	6 (14)	3 (7)	0 (0)	0 (0)	10 (15)	7 (10)
Extra work	3 (30)	3 (30)	14 (56)	13 (52)	1 (20)	0 (0)	18 (45)	16 (40)	12 (48)	15 (60)	18 (43)	24 (57)	0 (0)	0 (0)	30 (44)	39 (57)
Place of residence Mostly big cities	0 (0)	1 (10)	4 (16)	3 (12)	1 (20)	1 (20)	5 (13)	5 (13)	4 (16)	2 (8)	7 (17)	10 (24)	1 (100)	1 (100)	12 (18)	13 (19)
Higher education	4 (40)	3 (30)	3 (12)	2 (8)	3 (60)	3 (60)	10 (25)	8 (20)	2 (8)	6 (24)	12 (29)	9 (21)	1 (100)	0 (0)	15 (22)	16 (24)
Total number	10	10	25	25	5	5	40	40	25	25	42	42	1	1	68	68

D=Deceased twin; S=Surviving co-twin

Table 35. Physical inactivity and some sociologic variables in the female death discordant pairs according to earlier questionnaires. Absolute numbers and prevalence rates in % (parenthesis).

	IHD								not IHD							
	MZ		DZ		XZ		Tot.		MZ		DZ		XZ		Tot.	
	D	S	D	S	D	S	D	S	D	S	D	S	D	S	D	S
Hardly any exercise	2 (25)	0	1 (11)	0	-	-	3 (18)	0	2 (7)	1 (3)	5 (10)	7 (15)	0	0	7 (9)	8 (10)
Change of place of work $\geq 5$ times	1 (13)	0	0	0	-	-	1 (6)	0	2 (7)	2 (7)	1 (2)	2 (4)	0	0	3 (4)	4 (5)
Extra work	2 (25)	0	1 (11)	3 (33)	-	-	3 (18)	3 (18)	5 (17)	8 (27)	15 (31)	16 (33)	0	0	20 (25)	24 (30)
Place of residence Mostly big cities	0	0	0	0	-	-	0	0	7 (23)	4 (13)	9 (19)	10 (21)	0	0	15 (19)	15 (19)
Higher education	2 (25)	2 (25)	2 (22)	1 (11)	-	-	4 (24)	3 (18)	7 (23)	7 (23)	10 (21)	10 (21)	0	1 (50)	17 (21)	18 (23)
Total number	8	8	9	9			17	17	30	30	48	48	2	2	80	80

D=Deceased twin; S=Surviving co-twin

report  
than  
to 10  
(18 %  
of th  
extra  
rence  
disco  
death  
appli  
(abov  
much  
there  
among  
with  
not-I  
Comme  
The s  
possil  
variou  
cardia  
1971)  
diseas  
Tot  
likely  
increa  
in the  
(1970)  
often  
pairs.  
It  
the di  
associ

2028559372

reported slightly more often among the IHD death discordant male pairs than among the not-IHD death discordant pairs (20 % vs 18 % compared to 10 % vs 10 %). Among the IHD death discordant female twin pairs, 3 (18 %) of the deceased twins had reported hardly any exercise vs none of the surviving co-twins. With regard to change of place of work, extra work and place of residence, there are no substantial differences between the surviving and the deceased twins in the IHD death discordant male and female pairs. None of the female twins in the IHD death discordant pairs had mostly lived in big cities, whereas this applied to 19 % in the not-IHD group. With regard to higher education (above compulsory school), the proportion of educated subjects is much the same in the IHD and not-IHD male death discordant pairs but there are more marked differences within the various zygoty groups. Among the female IHD death discordant pairs the proportion of twins with higher education is 24 % vs 18 % compared to 21 % vs 23 % in the not-IHD group. None of the differences is statistically significant.

#### Comments

The sociologic data in the present study concern factors which may possibly predispose for IHD. However, precipitating factors, such as various life changes, have been shown to precede the onset of myocardial infarction (Theorell, 1971) and sudden death (Rahe & Lind, 1971) but the accumulation of life changes prior to the onset of disease does not seem to be a specific sign for IHD (Rahe, 1969).

To judge from the results of this study, it does not seem very likely that the social factors recorded could be responsible for the increased prevalence of IHD manifestations in the surviving co-twins in the IHD death discordant pairs. In the twin study by Liljefors (1970), ambition and overtime work were noted significantly more often among the affected partners in the infarction discordant MZ pairs.

It should be born in mind, however, that in retrospective studies the diseased subjects often tend to be more aware of signs and symptoms associated with the disease in question.

#### 4. COMBINATION OF FACTORS ASSOCIATED WITH ISCHAEMIC HEART DISEASE AMONG THE SURVIVING CO-TWINS

Prospective epidemiological studies have demonstrated the highest risk of developing IHD in subjects with multiple "risk factors" (Dawber et al. 1957; Kannel et al, 1961; Kannel et al, 1967; Tibblin & Wilhelmsen, 1971; Böttiger & Carlson, 1972; Stamler & Epstein, 1972). Data from the Pooling Project in the U.S.A. (Stamler & Epstein, 1972) show that the risk of death in IHD for white males aged 30-59 increased fivefold if they had three of the major risk factors (serum cholesterol  $\geq 250$  mg/dl; diastolic blood pressure  $> 90$  mm Hg; cigarette smoking). In the Stockholm Prospective Study, male subjects with 4 "risk factors" (any four of cholesterol, triglycerides, systolic blood pressure, diastolic blood pressure, hemoglobin and cigarette smoking) had a 10 times higher risk of developing a myocardial infarction.

Material: see Chapter II.

##### Methods

Before combining "risk factors" it is necessary to define "risk levels" for each factor. The methods for this are described in the presentation of the separate factors. The following "risk factors" and "risk levels" have been used:

- 1) Relative weight  $\geq 110$ .
- 2) Basal systolic blood pressure  $\geq 160$  mm Hg and/or basal diastolic blood pressure phase 4  $\geq 95$  mm Hg; or on antihypertensive treatment and not reaching the defined blood pressure levels.
- 3) Cholesterol  $\geq 250$  mg/100 ml and/or triglycerides  $\geq 150$  mg/100 ml; or on lipid reducing drugs and not reaching the defined lipid levels.
- 4) Diabetes mellitus: history of clinical diabetes mellitus.
- 5) Cigarette smoking: present cigarette smoking as elucidated from the mailed questionnaire 1967/1970.
- 6) Physical inactivity: hardly any exercise during 25-50 years of age as elucidated from the mailed questionnaire 1967/1970.
- 7) Extra work: Extra work  $\geq$  periodically beside the regular job as elucidated from the mailed questionnaire 1967/1970.

##### Results

The distribution of certain data of the four and visual IHD have on co-twins with partners died than the DZ the men who co-twins had but the results the same as

In Table Methods, and None of the noted for

The environment increased in male MZ co-2.8 "risk factor" proportion the pooled agreement in factors". Among whose partners factors, while DZ co-twins

Tables of biometric Of the male one biometric died from 0 are 94 % and the combination There are two co-twins with partners died figures are 73 % and 75 and 88 % and

## Results

The distribution of surviving co-twins according to the number of certain defined biometric "risk factors" (the possible combinations of the four factors defined under Methods) is presented in Table 36 and visualized in Fig. 8. The male co-twins whose partners died from IHD have on average 1.3 "risk factors" as compared to 1.2 among the co-twins whose partners died from other causes. Among the men whose partners died from IHD, the MZ co-twins have more "risk factors" than the DZ co-twins (1.6 vs 1.3) and the same trend applies among the men whose partners died from other causes (1.4 vs 1.1). The female co-twins have on average proportionally more biometric "risk factors" but the trends in the differences between zygosity groups are about the same as in the males.

In Table 37 and Fig. 9, three environmental factors, defined under Methods, are also included in the possible combination of "risk factors". None of the co-twins had all the "risk factors" but six of them were noted for one male MZ co-twin whose partner died from IHD.

The environmental factors contribute proportionally less to the increased number of risk factors per person among the females. The male MZ co-twins whose partners died from IHD now have on average 2.8 "risk factors" per person, mainly due to the relatively high proportion of present cigarette smokers. However, comparisons between the pooled male zygosity groups (IHD/not IHD) show fairly close agreement in prevalence rates throughout the different numbers of "risk factors". Among the women it is worth noting that the MZ co-twins, whose partners died from IHD, recorded only one of the environmental factors, which explains the reverse difference here between MZ and DZ co-twins within the IHD group.

Tables 38-39 display the cumulative distribution of the number of biometric as well as biometric plus environmental "risk factors". Of the male co-twins whose partners died from IHD, 80 % have at least one biometric risk factor as compared to 69 % of those whose partners died from other causes. The corresponding figures among the females are 94 % and 85 %. When environmental factors are also included in the combination, nearly all the twins have at least one "risk factor". There are two or more biometric "risk factors" in 40 % of the male co-twins with partners who died from IHD and in 38 % of those whose partners died from other causes. Among the females the corresponding figures are 82 % and 56 %. Including the environmental "risk factors", 73 % and 75 % of the male co-twins have at least two "risk factors", and 88 % and 71 % of the females.



Table 36. Distribution of respondent co-twins according to number of certain defined biometric "risk factors".

Number of factors	MALES								FEMALES							
	IHD				not IHD				IHD				not IHD			
	MZ	DZ	XZ	Tot.	MZ	DZ	XZ	Tot.	MZ	DZ	XZ	Tot.	MZ	DZ	XZ	Tot.
	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)
0	2 (20)	4 (16)	2 (40)	8 (20)	5 (20)	16 (38)	0 (0)	21 (31)	0 (0)	1 (11)	- (0)	1 (6)	4 (13)	8 (17)	0 (0)	12 (15)
1	3 (30)	11 (44)	2 (40)	16 (40)	10 (40)	11 (26)	0 (0)	21 (31)	1 (13)	1 (11)	- (0)	2 (12)	8 (27)	15 (31)	0 (0)	23 (29)
2	3 (30)	8 (32)	1 (20)	12 (30)	5 (20)	11 (26)	1 (100)	17 (25)	3 (38)	4 (44)	- (0)	7 (41)	9 (30)	14 (29)	0 (0)	23 (29)
3	1 (10)	2 (8)	0 (0)	3 (8)	4 (16)	4 (10)	0 (0)	8 (12)	4 (50)	3 (33)	- (0)	7 (41)	9 (30)	10 (21)	2 (100)	21 (26)
4	1 (10)	0 (0)	0 (0)	1 (3)	1 (4)	0 (0)	0 (0)	1 (1)	0 (0)	0 (0)	- (0)	0 (0)	0 (2)	1 (2)	0 (0)	1 (1)
Factors/co-twin	1.6	1.3	0.8	1.3	1.4	1.1	2.0	1.2	2.4	2.0	-	2.2	1.8	1.6	3.0	1.7
Numbers of co-twins	10	25	5	40	25	42	1	68	8	9	-	17	30	48	2	80

Table 37. Distribution of respondent co-twins according to number of certain defined biometric and environmental "risk factors".

Number of factors	MALES								FEMALES							
	IHD				not IHD				IHD				not IHD			
	MZ	DZ	XZ	Tot.	MZ	DZ	XZ	Tot.	MZ	DZ	XZ	Tot.	MZ	DZ	XZ	Tot.
	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)

2028559376

Table 37. Distribution of respondent co-twins according to number of certain defined biometric and environmental "risk factors".

Number of factors	MALES								FEMALES							
	IHD				not IHD				IHD				not IHD			
	MZ No. (%)	DZ No. (%)	XZ No. (%)	Tot. No. (%)	MZ No. (%)	DZ No. (%)	XZ No. (%)	Tot. No. (%)	MZ No. (%)	DZ No. (%)	XZ No. (%)	Tot. No. (%)	MZ No. (%)	DZ No. (%)	XZ No. (%)	Tot. No. (%)
0	0	0	1 (20)	1 (3)	0	3 (7)	0	3 (4)	0	0	-	0	4 (13)	2 (4)	0	6 (8)
1	2 (20)	7 (28)	1 (20)	10 (25)	3 (12)	11 (26)	0	14 (21)	1 (13)	1 (11)	-	2 (12)	3 (10)	14 (29)	0	17 (21)
2	4 (40)	7 (28)	3 (60)	14 (35)	12 (48)	14 (33)	0	26 (38)	3 (38)	3 (33)	-	6 (35)	8 (27)	12 (25)	0	20 (25)
3	1 (10)	8 (32)	0	9 (23)	6 (24)	9 (21)	1 (100)	16 (24)	3 (38)	3 (33)	-	6 (35)	11 (37)	12 (25)	2 (100)	25 (31)
4	1 (10)	2 (8)	0	3 (8)	3 (12)	4 (10)	0	7 (10)	1 (13)	2 (22)	-	3 (18)	3 (10)	7 (15)	0	10 (13)
5	1 (10)	1 (4)	0	2 (5)	1 (4)	1 (2)	0	2 (3)	0	0	-	0	1 (3)	1 (2)	0	2 (3)
6	1 (10)	0	0	1 (3)	0	0	0	0	0	0	-	0	0	0	0	0
Factors/co-twin	2.8	2.3	1.4	2.3	2.5	2.1	3.0	2.2	2.5	2.7	-	2.6	2.3	2.2	3.0	2.3
Number of co-twins	10	25	5	40	25	42	1	68	8	9	-	17	30	48	2	80

Table 38. Respondent co-twins tabulated according to the cumulative distribution of certain defined biometric "risk factors".

Number of "risk factors"	MALES								FEMALES							
	IHD				not IHD				IHD				not IHD			
	MZ	DZ	XZ	Tot.	MZ	DZ	XZ	Tot.	MZ	DZ	XZ	Tot.	MZ	DZ	XZ	Tot.
	No.	No.	No.	No.	No.	No.	No.	No.	No.	No.	No.	No.	No.	No.	No.	No.
	(%)	(%)	(%)	(%)	(%)	(%)	(%)	(%)	(%)	(%)	(%)	(%)	(%)	(%)	(%)	(%)
≥ 1	8	21	3	32	20	26	1	47	8	8	-	16	26	40	2	68
	(80)	(84)	(60)	(80)	(80)	(62)	(100)	(69)	(100)	(89)	-	(94)	(87)	(83)	(100)	(85)
≥ 2	5	10	1	16	10	15	1	26	7	7	-	14	18	25	2	45
	(50)	(40)	(20)	(40)	(40)	(36)	(100)	(38)	(88)	(78)	-	(82)	(60)	(52)	(100)	(56)
≥ 3	2	2	0	4	5	4	0	9	4	3	-	7	9	11	2	22
	(20)	(8)		(10)	(20)	(10)		(13)	(50)	(33)	-	(41)	(30)	(23)	(100)	(28)
≥ 4	1	0	0	1	1	0	0	1	0	0	-	0	0	1	0	1
	(10)			(3)	(4)			(1)			-			(2)		(1)
≥ 5	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0	0
Number of co-twins	10	25	5	40	25	42	1	68	8	9	-	17	30	48	2	80


Table 39. Respondent co-twins tabulated according to the cumulative distribution of certain defined biometric and environmental "risk factors".

Number of "risk	MALES								FEMALES							
	IHD				not IHD				IHD				not IHD			
	MZ	DZ	XZ	Tot.	MZ	DZ	XZ	Tot.	MZ	DZ	XZ	Tot.	MZ	DZ	XZ	Tot.

828559378

Table 39. Respondent co-twins tabulated according to the cumulative distribution of certain defined biometric and environmental "risk factors".

Number of "risk factors"	MALES								FEMALES							
	IHD				not IHD				IHD				not IHD			
	MZ No. (%)	DZ No. (%)	XZ No. (%)	Tot. No. (%)	MZ No. (%)	DZ No. (%)	XZ No. (%)	Tot. No. (%)	MZ No. (%)	DZ No. (%)	XZ No. (%)	Tot. No. (%)	MZ No. (%)	DZ No. (%)	XZ No. (%)	Tot. No. (%)
✓ 1	10 (100)	25 (100)	4 (80)	39 (98)	25 (100)	39 (93)	1 (100)	65 (96)	8 (100)	9 (100)	-	17 (100)	26 (87)	46 (96)	2 (100)	74 (93)
✓ 2	8 (80)	18 (72)	3 (60)	29 (73)	22 (88)	28 (67)	1 (100)	51 (75)	7 (88)	8 (89)	-	15 (88)	23 (77)	32 (67)	2 (100)	57 (71)
✓ 3	4 (40)	11 (44)	0	15 (38)	10 (40)	14 (33)	1 (100)	25 (37)	4 (50)	5 (56)	-	9 (53)	15 (50)	20 (42)	2 (100)	37 (46)
✓ 4	3 (30)	3 (12)	0	6 (15)	4 (16)	5 (12)	0	9 (13)	1 (13)	2 (22)	-	3 (18)	4 (13)	8 (17)	0	12 (15)
✓ 5	2 (20)	1 (4)	0	3 (8)	1 (4)	1 (2)	0	2 (3)	0	0	-	0	1 (3)	1 (2)	0	2 (3)
✓ 6	1 (10)	0	0	1 (3)	0	0	0	0	0	0	-	0	0	0	0	0
✓ 7	0	0	0	0	0	0	0	0	0	0	-	0	0	0	0	0
Number of co-twins	10	25	5	40	25	42	1	68	8	9	-	17	30	48	2	80




NUMBER OF "RISK FACTORS"


Fig. 8. Distribution of respondent co-twins according to certain defined biometric "risk factors". Black columns indicate that the partners died from IHD and white columns that the partners died from other causes.

Fig. 9. Distribution of respondent co-twins according to certain defined biometric "risk factors". Black columns indicate that the partners died from IHD and white columns that the partners died from other causes.

# Males



# Females



NUMBER OF "RISK FACTORS"

Fig. 9. Distribution of respondent co-twins according to certain defined biometric and environmental "risk factors". Black columns indicate that the partners died from IHD and white columns that the partners died from other causes.

None of the differences with regard to the number of risk factors is statistically significant.

### Comments

Before combining "risk factors" it is necessary to define "risk levels" for each factor. The cut-off points used here have been chosen fairly liberally, which naturally increases the sensitivity but decreases the specificity with regard to the risk of IHD. It seems, however, that the risk rises steadily from the lowest to the highest values for most of the continuous risk factors (Dawber & Kannel, 1972; Stamler & Epstein, 1972).

New statistical methods for multivariate analyses have been applied more recently to evaluate the risk of IHD when several risk characteristics are considered simultaneously. In the Framingham study, a multiple logistic model was used to compute risk probabilities for each individual (Truett et al, 1967). Coefficients for each of the risk factors were based on the risk factor level at the first examination and the 12-year IHD incidence.

The relative importance of risk factors is, however, not the same in all populations. Furthermore, the multivariate risk functions have not yet been translated into clinical terms. A combination of risk factors defined by arbitrarily chosen levels can therefore be useful as a broad classification of high risk individuals.

In the present study, the number of risk factors does not differ significantly between co-twins whose partners died from IHD and those whose partners died from other causes. When only biometric risk factors were combined, the same tendencies are found for both males and females with more risk factors among the MZ than the DZ co-twins, irrespective of the cause of death of the partner (IHD/notIHD). Furthermore, the co-twins with partners who died from IHD have somewhat more risk factors than those whose partners died from other causes, the difference being more pronounced among the females.

Twin studies cited in connection with the separate risk factors have shown that the biometric risk factors are governed to varying degrees by heredity but may also be influenced by the environment. In a prospective study of 50-year-old men in Gothenburg, the occurrence of risk factors had a predictive value not only for the incidence of myocardial infarction but also for early death from other causes (Tibblin, 1972). The differences in number of risk factors between MZ and DZ co-twins irrespective of the cause of death of the partner might thus reflect constitutional differences. The male co-twins with partner

who died from  
whose parts  
marked, possibly  
for IID as  
factor such  
number of  
fraction ar

The diff  
same trends  
physical in  
combination  
tionally le  
especially  
the males.  
found the o  
determined  
sidered env  
of IHD in ma

In a con  
with respect  
concluded th  
only part of  
and women. S  
however, mor

The pres-  
 significantly dis-  
 from LHD and  
 could be due  
 cles, howeve  
 factors amon  
 differences  
 tutions but  
 dently of th  
 Firmer concl  
 follow-up of

who died from IHD have rather more biometric risk factors than those whose partners died from other causes but the difference is not very marked, possibly because the so-called risk factors are not as specific for IHD as used to be thought. According to Tibblin (1972), some common factor such as premature aging could be responsible for an increased number of risk factors as well as the development of myocardial infarction and death from non-cardiovascular causes.

The differences in the number of biometric risk factors show the same trends when environmental risk factors (cigarette smoking, physical inactivity and extra work) are included in the possible combination. The environmental factors, however, contributed proportionally less to the increased number of factors among females, especially the MZ co-twins whose partners died from IHD, than among the males. Harvald & Hauge (1968), in their study on Danish twin pairs, found the occurrence of fatal "coronary occlusion" to be genetically determined to a much larger extent in females than males. They considered environmental factors to play a greater part in the etiology of IHD in males compared to females.

In a comparison between male and female populations in Gothenburg with respect to risk factors for IHD (Bengtsson et al, 1973), it was concluded that sex differences in these factors apparently explain only part of the different incidences of myocardial infarction in men and women. Smoking, alcohol consumption and stress experience were, however, more common among the males.

The present differences in the number of risk factors do not significantly discriminate the surviving co-twins with partners who died from IHD and those whose partners died from other reasons, which partly could be due to a lack of specificity in the risk factors. The tendencies, however, are consistent with a somewhat higher number of risk factors among the co-twins whose partners died from IHD. The slight differences in the number of risk factors may reflect different constitutions but it is also probable that genetic factors operate independently of the established risk factors in the development of IHD. Firmer conclusions in this respect will have to await a mortality follow-up of the examined co-twins.



#### IV GENERAL DISCUSSION

Epidemiological studies have indicated that IHD has a multifactorial etiology. From the standpoint of preventive action, most interest has been focused on the environmental factors, which can be modified and controlled, and there have been relatively few studies on the importance of familial factors and the genetic predisposition for IHD, which seem to be especially important for detecting individuals at highest risk. Twin studies probably offer the best tool with which to study the relative influence of genetic and environmental factors.

Various theoretical models in twin studies were discussed at an international twin symposium in Puerto Rico in 1969 but it was also pointed out that there is no generally accepted methodology in the design of twin studies and that the possibilities are many and complex. It was suggested that clinical examination of the partners of twins who have died could be considered as a valuable extension of mortality studies.

One of the main purposes of the present study is to evaluate a genetic influence in IHD - by examining whether the occurrence of IHD in the surviving co-twin is associated with the partner's cause of death (IHD/not IHD).

The present material of death discordant pairs comprises 78.2 % of a sample selected from the Swedish Twin Registry. This response rate must be considered fairly high, partly because the population was drawn from the whole of Sweden and partly in view of the fact that the co-twins had lost their partners on average 5 months earlier.

The accuracy of cause of death determinations depends mainly on a high autopsy rate. Because the autopsy rate for the deceased twins in the present material was about 57 %, other sources of information (hospital records etc.) were consulted in order to make the mortality evaluation as reliable as possible.

This was especially important because the IHD material included cases of sudden death (ISC number 795.99). If there was a strong suspicion that a case of sudden death had some other explanation than IHD as the underlying cause, then it was not included in the IHD group. About 70 % of the sudden deaths included as IHD cases had either a preexisting history of IHD (myocardial infarction and/or angina pectoris on effort) or findings at autopsy indicating IHD (signs of fresh or old infarct and/or severe atherosclerotic changes of the coronary arteries) as compared to about 4 % of the twins who died from other causes than

IHD. The other evidence or other reason other studies was found to et al, 1966; of the occurrence twins have I (not IHD). In discordant myocardial in many epid because about IHD but also in general a et al, 1973) discordant p

The occurrence by determination and ECG infarction was partners died causes. Among clearest with the difference valence of m DZ co-twins m According to represents on depressions i specific pred The appearance been shown to Åstrand & Lund Using ST depr among the sur the female co higher than as In the males but in the fem These differer impression of

IHD. The other 30 % of the IHD sudden deaths had no such earlier evidence of this disorder; they were included because there was no other reasonable explanation for the death. This is in line with other studies, in which the initial clinical manifestation of IHD was found to be sudden death in about 20-25 % of IHD cases (Kuller et al, 1966; Kuller, 1966). In the present investigation, analyses of the occurrence of IHD and risk factors among the surviving co-twins have been related to the cause of death of the partner (IHD, not IHD). In addition to the cases of sudden death, the IHD death discordant pairs thus include those where the cause of death was myocardial infarction. Pooling these two diagnoses, as has been done in many epidemiological studies, seems to be justifiable not only because about 70 % of the sudden deaths showed evidence of previous IHD but also because the risk factors for the sudden death cases are in general agreement with those for myocardial infarction (Vedin et al, 1973). Neither did the relatively small number of IHD death discordant pairs permit a further breakdown into subgroups.

The occurrence of IHD in the surviving co-twins was established by determining the prevalence of angina pectoris, myocardial infarction and ECG changes suggestive of IHD. Angina pectoris and myocardial infarction was found more commonly among the surviving co-twins whose partners died from IHD than among those whose partners died from other causes. Among females this was true for angina pectoris. The trend was clearest with regard to myocardial infarction among males but none of the differences is statistically significant. The relatively low prevalence of manifest IHD in conjunction with the small numbers of MZ and DZ co-twins makes it difficult to assess the significance of heredity. According to Epstein (1964), a prevalence rate of 5.5 % for manifest IHD represents only the "top of the iceberg". It has been considered that ST depressions in connection with exercise could be a fairly sensitive and specific predictor of coronary obstructive disease (Helfant et al, 1973). The appearance of ST segmental depressions in response to exercise has been shown to be a powerful predictor of overt IHD (Mattingly, 1962; Åstrand & Lundman, 1968; Blackburn et al, 1970; Doyle & Kinch, 1970). Using ST depressions in connection with exercise as a sign of IHD among the surviving co-twins, the frequency among both the male and the female co-twins whose partners died from IHD was significantly higher than among the co-twins whose partners died from other causes. In the males the difference was significant for ST depressions  $\geq 1$  mm but in the females only when ST depressions of  $\geq 0.5$  mm were included. These differences certainly reflect constitutional differences. The impression of a substantial genetic influence in IHD is strengthened

by the finding that the male MZ co-twins whose partners died from IHD displayed significantly more IHD manifestations than those MZ co-twins whose partners died from other causes despite the fact that the group with partners who died from IHD was about 5 years younger.

It is well known that ST depressions seem to be a less specific sign of IHD in women (Biörck, 1946; Lepeschkin, 1958; Åstrand, 1965). They were more common among the females than among the males in the present study and did not discriminate the genetically predisposed individuals from the others as effectively as among the males. It has been conjectured that ECG changes suggestive of IHD in women may reflect hypertension rather than IHD (Bengtsson, 1973). It is thus conceivable that the relatively high blood pressure, especially the casual, in the female co-twins whose partners died from IHD, accounts to some extent for the difference in IHD manifestations. This possibility is strengthened by the fact that if one excludes ST depressions accompanied by high R-wave amplitudes (code 3:1), which are thought to reflect a left ventricular hypertrophy due to hypertension, then the differences in the females are no longer significant. It must, however, be born in mind that the level of blood pressure has been shown by some authors (Takkunen, 1964; Lundman, 1966) to be under a relatively strong genetic influence, so that the differences with regard to IHD manifestations in the females could also reflect constitutional differences. In the Framingham study, electrocardiographic evidence of left ventricular hypertrophy was strongly associated with both hypertension and IHD; when the criteria for left ventricular hypertrophy included ST depression and T-wave inversion, IHD incidence was predicted over and above what could be accounted for by the concurrent hypertension (Kannel et al, 1970; Dawber & Kannel, 1972). It was suggested that these abnormalities reflect, in addition to hypertensive hypertrophy, an ischaemic myocardial involvement due to associated coronary atherosclerosis.

One of the other purposes of the present investigation was to study whether there is an association between the risk factor profile of the surviving co-twin and the partner's cause of death (IHD/not IHD). For this purpose the factors found to be associated with IHD were divided into biometric and environmental. It is probable that the determinants of many of the biometric factors are mostly genetic but they may be susceptible to environmental influence, too. Similarly environmental factors may be linked in some measure with the genetic constitution (Biörck, 1959).

For most of the separate biometric risk factors recorded, the values were consistently somewhat higher among both the male and the

female co-partners of MZ co-twins. They differ somewhat from the rest of the twins who were on average the same in causes that

Further, 4 years older. Some of the twins is thus possibly somewhat younger. However, about the same for females, though about 4 years older. They died from causes found with partners who are to some extent

Concerning information, twins in the study at that time, ~~perhaps~~ seemed advised because the surviving co-twin about the death

With regard to and some similarity between the pairs. The proportion of the not-IHD smokers was 60% in pairs (60% in

Most especially in 1964; Best, No association born in mind and XZ male

female co-twins whose partners died from IHD, compared to those whose partners died from other causes and this was especially true for the MZ co-twins. It must be remembered, however, that the zygosity groups differ somewhat in age, which could possibly affect the interpretation of the results. The male MZ co-twins whose partners died from IHD were on average nearly 5 years younger than both the DZ co-twins in the same IHD group and the MZ co-twins whose partners died from other causes than IHD.

Furthermore, the male DZ co-twins with IHD partners were about 4 years older than those whose partners had not died from IHD. As some of the biometric risk factors are more or less age dependent, it is thus possible that the differences between the MZ co-twins are somewhat underestimated, whereas the reverse is true for the DZ co-twins. However, the pooled male zygosity groups (IHD/not IHD) had about the same mean age. There was also an age difference among the females, the MZ and DZ co-twins whose partners died from IHD being about 4 years older than the corresponding co-twins whose partners died from other reasons. It is thus possible that the differences found with respect to, for instance, blood pressure and cholesterol are to some extent due to the difference in age.

Concerning the environmental factors, except alcohol registration, information was obtained from questionnaires mailed to all twins in the Swedish Twin registry in the years 1967 and 1970. At that time, all the present death discordant pairs were unbroken. It seemed advisable to use the information in earlier questionnaires because the death of the partner might have distorted answers from surviving co-twins as well as those obtained through relatives about the deceased partners.

With regard to smoking, alcohol registration, physical inactivity and some social factors, no substantial differences were found between the deceased and the surviving twins in the death discordant pairs. The IHD death discordant pairs included a somewhat higher proportion of total smokers as well as cigarette smokers, compared to the not-IHD death discordant pairs, and the proportion of cigarette smokers was especially high among the male IHD death discordant MZ pairs (60 % among the deceased partners and 80 % among the survivors).

Most epidemiological studies point to a positive relation, especially in men, between mortality in IHD and smoking (Doll & Hill, 1964; Best, 1966; Hammond, 1966; Kannel, 1966; Fletcher & Horn, 1970). No association was found in the present investigation but it must be born in mind that the numbers involved are small. The surviving MZ and XZ male co-twins whose partners died from IHD were cigarette

smokers, especially  $\geq 11$  cigarettes daily, proportionally more often than the other male MZ and XZ twins, a circumstance which may have contributed to the corresponding difference in IHD manifestations. However, the deceased MZ and XZ twins also displayed the same tendencies in the smoking of cigarettes. Thus it is possible that the differences with regard to cigarette smoking merely reflect differences in constitution, which could play the major part also with regard to the development of IHD.

Among the men born in 1913 (Tibblin, 1972), it seemed that being registered with the local temperance board was strongly associated with myocardial infarction. All the male death discordant twins in the present investigation have been matched against a register for misconduct involving alcohol. Registration was somewhat more common, although not significantly so, among the deceased twins, irrespective of the cause of death (IHD/not IHD).

The questionnaire data used in this study on sociologic background variables represent factors which could predispose for IHD. These findings, however, do not preclude the operation of precipitating factors such as various life changes, which have been shown to precede the onset of myocardial infarction (Theorell, 1971) and sudden death (Rahe & Lind, 1971).

The highest risk of developing IHD has been demonstrated among individuals with multiple "risk factors" (Dawber et al, 1957; Kannel et al, 1961; Kannel et al, 1967; Tibblin & Wilhelmsen, 1971; Böttiger & Carlson, 1972; Stamler & Epstein, 1972). The risk factors used in the combination were defined with fairly liberal cut-off points, which is a practical and common way of defining risk levels. Risk evaluation by means of new statistical methods for multivariate analyses seems to be a more physiological approach, considering that the risk steadily increases from the lowest to the highest values for most of the continuous variables (Dawber & Kannel, 1972; Stamler & Epstein, 1972). These methods have not yet been translated into clinical terms and the coefficients for each of the risk factors are based upon special populations. Using a multivariate risk function (Wilhelmsen et al, 1973) based on the variables cholesterol, systolic blood pressure and smoking, it has been shown (Elmfeldt, 1974) that a fairly high proportion of men developed myocardial infarction notwithstanding their classification as "low risks".

When only biometric risk factors were combined in the present study, more risk factors were noted among the MZ co-twins than the DZ co-twins, irrespective of the cause of death (IHD/not IHD). The same trend was found when the environmental risk factors (cigarette

smoking, possible tionally the female than among a less im Hauge, 19 of risk f twins who from other cient spe spective most of ti specific appear to

To ju that a cor ment of II establishe or a gener low "life underlying have propo IHD could important not been id Epstein (10 disposing metabolic are more c show more more concl of these t instituted

smoking, physical inactivity and extra work) were included in the possible combination. The environmental factors contributed proportionally less to the increase in the number of risk factors among the females, especially the MZ co-twins whose partners died from IHD, than among the males. Environmental factors have been found to play a less important role in the development of IHD in women (Harvald & Hauge, 1968; Bengtsson, 1973). The differences found in the number of risk factors did not significantly discriminate the surviving co-twins whose partners died from IHD from those whose partners died from other reasons, a circumstance which could be due to an insufficient specificity of the risk factors. Incidence data from the prospective study on men born in 1913 (Tibblin, 1972) have shown that most of the risk factors which have been considered to be rather specific for the subsequent occurrence of myocardial infarction also appear to be almost as valid for early death from whatever cause.

To judge from the results of the present study, it can be concluded that a considerable genetic influence seems to operate in the development of IHD. Genetic influences are transmitted through some of the established risk factors, but it is possible that unknown risk factors or a general biologic factor as "premature ageing" (Tibblin, 1972) or low "life potential" (Björck, 1974) constitute the most important underlying factor in the development of IHD. Dawber & Kannel (1972) have proposed, that the continued high morbidity and mortality from IHD could be a consequence of ageing and genetic make-up and that the important factors contributing to the development of this disease have not been identified satisfactorily. It has also been proposed by Epstein (1964) that if one could identify and measure all of the predisposing traits as underlying biologic disturbances in terms of metabolic or other defects, then it would probably emerge that they are more common than the prevalence of the disease would suggest and show more clear-cut distributions among family members. It was furthermore concluded that the prevention of IHD demands that the carriers of these traits be identified, so that prophylactic measures can be instituted at an early age among genetically susceptible individuals.

## GENERAL SUMMARY AND CONCLUSIONS

All the twins in the Swedish Twin Registry have been the subject of a continuous mortality follow-up ever since the Registry was established in 1959-1961. The complete twin registry is matched regularly against a total death registry for Sweden at the Central Bureau of Statistics and this procedure makes it possible to obtain the death certificates. Since 1971 the matching has been done every month. This procedure was a prerequisite for the present study.

The principal object of this investigation was to evaluate the genetic influence in IHD in the examined co-twins and to create a basis for their further mortality follow-up, thereby also paving the way for assessments of the predictive value of the measured risk factors and the hereditary influence.

All unbroken male and female twin pairs from the Swedish Twin Registry below the age of 70 who, in the period January 1st, 1971 to March 15th, 1973, became death discordant, were selected for the study. 78.2% or 205 of the surviving co-twins could be examined on average about 5 months after the death of the partner.

The cause of death of the partner was established by a team of physicians on the basis of all the assembled information (hospital records, autopsy protocols etc.). The causes of death were classified according to the International Statistical Classification of Diseases.

The zygosity diagnoses were taken from the record in the Twin Registry, which is based upon questions as to similarity. If the twins in a pair had given conflicting answers, they were classified as unknown zygosity (XZ).

Among 108 male death discordant pairs, the cause of death was IHD in 40 pairs (10 MZ, 25 DZ and 5 XZ) and other than IHD in 68 pairs (25 MZ, 42 DZ and 1 XZ). Among 97 female death discordant pairs, the cause of death was IHD in 17 pairs (8 MZ, 9 DZ) and other than IHD in 80 pairs (30 MZ, 48 DZ and 2 XZ).

191 of the surviving co-twins were examined at the Seraphimer Hospital and the other 14 at their local hospital.

A sociologic and medical history was taken, using questionnaires. The diagnosis of angina pectoris was established by interview according to the questionnaire designed by the London School of Hygiene and Tropical Medicine. Myocardial infarction was considered established when it had been verified at hospital. Blood pressure determinations, anthropometric measurements and X-ray of the heart and lungs were

performed  
Blood sample  
cholesterol  
sedimentat  
The urine

Myocard  
co-twins w  
twins and  
the surviv  
than IHD.  
tion. If as  
sions  $\geq 0.5$   
criteria of  
whose part  
( $< 0.05$ ) mo  
In the male  
even when  
Similarly,  
significant  
 $\geq 1.0$  mm, as  
whose part  
in the pres  
are thought  
tension, th

Most of  
thickness,  
values for  
whose part  
blood press  
diabetes me  
discordant  
found with  
tion rate, v  
IHD. Inform  
smoking hab  
work, educat  
criminate th  
cordant pair  
85 %) compar  
daily was si  
male MZ co-t  
partners die

performed as well as ECG before, during and after an ergometer test. Blood samples were drawn after an overnight's fast for analyses of cholesterol, triglycerides, uric acid and blood sugar. Erythrocyte sedimentation rate, hemoglobin and hematocrit were also determined. The urine was tested for proteinuria and glucosuria.

Myocardial infarction had occurred in 15 % of the surviving male co-twins whose partners had died from IHD (3 or 30 % of the MZ co-twins and 3 or 12 % of the DZ co-twins) as compared to only 3 % of the surviving co-twins whose partners had died from other causes than IHD. Only one of the female co-twins had had myocardial infarction. If angina pectoris, pathologic Q-wave and segmental ST depressions  $\geq 0.5$  mm in connection with exercise were also included in the criteria of IHD, both male and female co-twins (pooled zygosity groups) whose partners died from IHD displayed these signs significantly ( $p < 0.05$ ) more often than those whose partners died from other causes. In the males, but not in the females, the difference was significant even when ST depressions  $\geq 1.0$  mm was included in the criteria of IHD. Similarly, the male MZ co-twins whose partners died from IHD displayed significantly more IHD manifestations ( $p < 0.05$  with ST depressions  $\geq 1.0$  mm, and  $p < 0.01$  with ST depressions  $\geq 0.5$  mm) than the MZ co-twins whose partners had died from other causes. Excluding ST depressions coded in the presence of high R-wave amplitudes (Minnesota code 3:1), which are thought to reflect a left ventricular hypertrophy due to hypertension, the differences in the females are no longer significant.

Most of the biometric factors measured (relative weight, skinfold thickness, blood pressure, lipids, uric acid) showed somewhat higher values for the co-twins whose partners died from IHD compared to those whose partners died for other reasons. The difference in casual systolic blood pressure was significant ( $p < 0.05$ ) among the females. Overt diabetes mellitus showed about the same prevalence in the two death discordant groups (IHD, not IHD). Nor were significant differences found with regard to hematocrit, hemoglobin or erythrocyte sedimentation rate, which some authors have designated as "risk factors" for IHD. Information obtained earlier through mailed questionnaires about smoking habits, physical inactivity, extra work, change of place of work, education and place of residence, did not significantly discriminate the deceased and the surviving co-twins. The IHD death discordant pairs had a somewhat higher proportion of smokers (83 % vs 85 %) compared to the not-IHD (76 % vs 72 %). To smoke  $\geq 11$  cigarettes daily was significantly ( $p < 0.05$ ) more common among the surviving male MZ co-twins whose partners died from IHD than among those whose partners died from other causes. The corresponding comparison between



the deceased male MZ co-twins gave the same trend but not so pronounced.

All male death discordant twins were matched against a nationwide register for misconduct while under the influence of alcohol. Registration was more common among the not-IHD death discordant pairs. The deceased partners had been registered somewhat more often, irrespective of the cause of death (IHD, not IHD).

The biometric "risk factors" (relative weight  $\geq 110$  according to a height-weight index, basal systolic blood pressure  $\geq 160$  mm Hg and/or basal diastolic blood pressure  $\geq 95$  mm Hg, cholesterol  $\geq 250$  mg/100 ml and/or triglycerides  $\geq 150$  mg/100 ml, and overt diabetes mellitus) were then combined with the environmental "risk factors" (cigarette smoking, extra work and physical inactivity). It was found that the surviving co-twins whose partners died from IHD had on average a somewhat higher number of biometric risk factors than those whose partners died from other causes, the difference being more pronounced among the female co-twins. The inclusion of environmental "risk factors" contributed proportionally less to the number of "risk factors" among the females, especially the MZ co-twins.

The conclusions from the present investigation could be summarized as follows:

- 1) The present results indicate a substantial genetic influence in the development of IHD.
- 2) In males the risk factors measured, singly and in combination, seem to explain only part of the difference in IHD manifestations between the surviving co-twins whose partners died from IHD and those, whose partners died from other causes. In females, however, elevated blood pressure may explain a great deal of the difference found in IHD manifestations. The genetic influences are probably transmitted not only through some of the risk factors measured but also through factors which are still unknown.
- 3) The environmental factors recorded by earlier questionnaires did not discriminate significantly between the deceased and the surviving co-twins.

I wish to  
study not

Profes  
Karolinska  
me and pla  
vided valu  
indepted b  
Environmen  
National H  
and suppor  
throughout

Torbje  
assisted i  
in the mor  
manuscript

Mr. Ru  
methodolog

Profess  
Professor  
M.D., for  
Göran Skog

Mr. Sta  
Lars Holml  
analyses.

Profess  
script.

Miss Ul  
death cert  
nent inform

Part of  
thanks are  
Carlstén, M.  
Sahlgrenska

Valuable  
by Miss Elsa

Miss Eva Bon  
Miss Bri

She also typ  
also due to

#### ACKNOWLEDGEMENTS

I wish to express my sincere gratitude to all who contributed to this study not forgetting all the co-operative twins.

Professor Gunnar Björck, M.D., Head of the Department of Medicine, Karolinska Institutet at Serafimerlasarettet, who not only supported me and placed the necessary facilities at my disposal but also provided valuable advice and stimulating points of view. I am also greatly indebted to Professor Lars Friberg, M.D., Head of the Department of Environmental Hygiene at the Karolinska Institutet and the Swedish National Environmental and Protection Board, for his valuable advice and support. He also took active part in the mortality evaluations throughout the study.

Torbjörn Lundman, M.D., who aroused my interest in twin methodology, assisted in the planning of the study, advised me and took active part in the mortality evaluations throughout the study. He also revised the manuscript.

Mr. Rune Cederlöf, Ph.D., who gave valuable advice and help in methodological problems.

Professor Bengt Pernow, M.D., Associate Professor John Wahren, M.D., Professor Rolf Blomstrand, M.D., and Associate Professor Gunnar Westberg, M.D., for placing personnel and laboratory facilities at my disposal. Göran Skogsberg, M.D., kindly examined all the X-rays.

Mr. Staffan Lundquist, fil.lic., Mr. Rein Maasing, fil.kand., and Mr. Lars Holmlund for competent assistance with the statistics and computer analyses.

Professor Jan Lindsten, M.D., for valuable comments on the manuscript.

Miss Ulla Lörich, fil. kand., who administered the collection of death certificates, hospital records, autopsy records and other pertinent information on the deceased twins.

Part of the examinations were performed at other hospitals and my thanks are due to a number of colleagues especially Professor Arne Carlstén, M.D., Head of the Department of Clinical Physiology, Sahlgrenska Hospital, Gothenburg.

Valuable assistance in the clinical examinations has been provided by Miss Elsa Munther, Mrs. Ingrid Ollén, Miss Anna Stina Hedén and Miss Eva Bonde.

Miss Britt Rosén for excellent and devoted secretarial assistance. She also typed the final version for offset printing. My thanks are also due to Miss Elisabeth Gyllenstierna, Mrs. Gunilla Haag and Mrs.

Birgitta Öhman for secretarial help and to Miss Ann-Marie Lothigius, for preparing the figures and to Mrs. Kim Hellman for enthusiastic help with the bibliography.

Mr. Patrick Hort, M.A., for careful correction of the English text.

The investigation was supported by grants and a Scholarship from the Swedish National Association against Heart and Chest Diseases, and also by grants from the Swedish Medical Research Council and the American Medical Association, Education and Research Foundation, for which I am most grateful.

1. ADELSON
2. ALLEN
3. ANTON
4. APD, A.  
Acta Med
5. ATHENS  
1970.
6. AUERBACH  
arteries
7. BAINES  
younger
8. BEAUMONT  
Classification
9. BENEDICT
10. BENGTS  
women
11. BERNHARDT  
Glucose  
infarction
12. BERNASCO  
50:176, 1
13. BEST, E.  
1966
14. BJÖRCK
15. BJÖRCK  
in a normal
16. BJÖRCK  
Infarction
17. BJÖRCK
18. BJÖRCK
19. BJÖRCK  
and degeneration
20. BJÖRCK  
Sverige
21. BJÖRCK
22. BJÖRCK  
Epidemiology
23. BJURULF
24. BJURULF  
disease
25. BJÖRNTORP  
human
26. BLACKBURN  
population
27. BLACKBURN  
cardiogram  
nine, Cardio
28. BLONQVIST  
1963
29. BLONQVIST
30. BLONQVIST  
sclerotic

## REFERENCES

1. ADELSON, L.: Sudden death from coronary disease - the cardiac conundrum, *Postgrad. Med. J.* 30:139, 1961.
2. ALLEN, G., HARVALD, B. & SHIELDS, J.: Measures of twin concordance, *Acta Genet. (Basel)*, 17:475, 1967.
3. ANTONOWSKY, A.: Social class and the major cardiovascular diseases, *J. Chronic Dis.* 21:65, 1968.
4. ARO, A.: Serum lipids and lipoproteins in first degree relatives of young survivors of myocardial infarction, *Acta Med. Scand. suppl.* 553, 1973.
5. ATHEROSCLEROSIS STUDY GROUP: Primary prevention of the atherosclerotic diseases, *Circulation* 42, suppl. 3, 1970.
6. AUERBACH, O., HAMMOND, E.C. & GARFINKEL, L.: Smoking in relation to atherosclerosis of the coronary arteries, *N. Engl. J. Med.* 273:775, 1965.
7. BAINTON, C.R. & PETERSON, D.R.: Deaths from coronary heart disease in persons fifty years of age and younger, A community-wide study, *N. Engl. J. Med.* 268:569, 1963.
8. BEAUMONT, J.L., CARLSON, L.A., COOPER, G.R., FEJFAR, Z., FREDRICKSON, D.S. & STRASSER, T.: Classification of hyperlipidaemias and hyperlipoproteinaemias, *Bull. WHO* 43, 1970, p. 891.
9. BENEDICT, R.B.: Coronary heart disease in identical female twins, *Am. J. Med.* 24:614, 1958.
10. BENGTISSON, C.: Ischaemic heart disease in women, A study based on a randomized population sample of women and women with myocardial infarction in Göteborg, Sweden, *Acta Med. Scand. suppl.* 549, 1973.
11. BERCHTOLD, P., BJÖRNTORP, P., GUSTAFSON, A., LINDHOLM, B., TIBBLIN, G. & WILHELMSON, L.: Glucose tolerance, plasma insulin and lipids in relation to adipose tissue cellularity in men after myocardial infarction, *Acta Med. Scand.* 191:35, 1972.
12. BERNASCONI, P., BROCHIER, M. & ZERAFKA, P.: Angine de poitrine chez deux jumeaux, *Arch. Mal. Coeur*, 50:176, 1957.
13. BEST, E.W.R.: A Canadian study of smoking and health, Department of National Health and Welfare, Ottawa, 1966.
14. BJÖRCK, G.: Hypoxaemia tests in coronary disease, *Br. Heart J.* 8:17, 1946.
15. BJÖRCK, G., BLOMQVIST, G. & SIEVERS, J.: Cholesterol values in patients with myocardial infarction and in a normal control group, *Acta Med. Scand.* 156:493, 1957.
16. BJÖRCK, G., BLOMQVIST, G. & SIEVERS, J.: Studies on myocardial infarction in Malmö 1935-1954. II. Infarction rate by occupational group, *Acta Med. Scand.* 161:21, 1958.
17. BJÖRCK, G.: Environment and disorders of the circulatory system, *Bull. N.Y. Acad. Med.* 35:3, 1959.
18. BJÖRCK, G.: Factors of risk in coronary artery disease, *Mal. Cardiov.* 4:41, 1963.
19. BJÖRCK, G. & BYLIN, G.: Comparative trends in four Scandinavian Countries of mortality from atherosclerotic and degenerative heart disease during the years 1952-1960, *Acta Med. Scand.* 177:765, 1965.
20. BJÖRCK, G., BOLANDER, A.M. & GYLLENHAMMAR, K.: Synpunkter på dödsfall i ischemisk hjärtsjukdom i Sverige, *Läkartidningen* 67:3993, 1970.
21. BJÖRCK, G. & WIKLAND, B.: "Sudden death" - what are we talking about? *Circulation* 45:256, 1972.
22. BJÖRCK, G.: Ideas and Concepts in the Prevention of Ischemic Heart Disease, To be publ. in *Reviews of Epidemiology and Social Medicine*, 1974.
23. BJURULF, P.: Atherosclerosis and body-build, *Acta Med. Scand. suppl.* 349, 1959.
24. BJURULF, P., GARLIND, T. & STERNBY, N.H.: On epidemiologic methods for recording ischaemic heart disease, *Acta Med. Scand. suppl.* 474, 1967.
25. BJÖRNTORP, P. & SJÖSTRÖM, L.: Number and size of adipose tissue fat cells in relation to metabolism in human obesity, *Metabolism* 20:703, 1971.
26. BLACKBURN, H., KEYS, A., SIMONSON, E., PAUTAHARJU, P. & PUNSAR, S.: The electrocardiogram in population studies, A classification system, *Circulation* 21:1160, 1960.
27. BLACKBURN, H., TAYLOR, H.L. & KEYS, A.: Coronary heart disease in seven countries, XVI. The electrocardiogram in prediction of five-year coronary heart disease incidence among men aged forty through fifty-nine, *Circulation* 41, suppl. 1, 154, 1970.
28. BLUMQVIST, G. & BJÖRCK, G.: Coronary mortality in relation to total mortality, *Acta Med. Scand.* 173:229, 1963.
29. BLUMQVIST, G.: The Frank lead/exercise electrocardiogram, *Acta Med. Scand. suppl.* 440, 1965.
30. BLUMQVIST, G.: Use of exercise testing for diagnostic and functional evaluation of patients with arteriosclerotic heart disease, *Circulation* 44:1120, 1971.

31. BOLANDER, A.-M.: Cause of death statistics - a tool in medical research and especially in epidemiological survey. Solicited paper to the 1973 General Conference of the International Union for the Scientific Study of Population, Statistiska Centralbyrån, Stockholm, 1973.
32. BRITTON, M.: Diagnostic errors discovered at autopsy. *Acta Med. Scand.* 1974. In Press.
33. BÖTTIGER, L.E. & CARLSON, L.A.: Stockholm Prospective study II - nio års uppföljning. *Läkartidningen* 69: 4640, 1972.
34. BÖTTIGER, L.E. & CARLSON, L.A.: Hemoglobin, plasma lipids, and coronary heart disease. *Am. Heart J.* 86:842, 1973.
35. BÖTTIGER, L.E., CARLSON, L.A., EKELUND, L.-G. & OLSSON, A.G.: Raised erythrocyte sedimentation rate in asymptomatic hyperlipidaemia. *Brit. Med. J.* 2:681, 1973.
36. BÖTTIGER, L.E.: Erythrocyte sedimentation rate and plasma lipids. *Acta Med. Scand.* 193:53, 1973.
37. CARLSON, L.A.: Serum lipids in normal men. *Acta Med. Scand.* 167:377, 1960.
38. CARLSON, L.A.: Serum lipids in men with myocardial infarction. *Acta Med. Scand.* 167:399, 1960.
39. CARLSON, L.A. & BÖTTIGER, L.E.: Ischaemic heart-disease in relation to fasting values of plasma triglycerides and cholesterol. Stockholm Prospective Study. *Lancet* 1:865, 1972.
40. CEDERLÖF, R., FRIBERG, L., JONSSON, E. & KAU, L.: Studies on similarity diagnosis in twins with the aid of mailed questionnaires. *Acta Genet.* 11:338, 1961.
41. CEDERLÖF, R.: The Twin Method in Epidemiological Studies on Chronic Disease, Karolinska Institutet, 1966 a.
42. CEDERLÖF, R., JONSSON, E. & KAU, L.: Respiratory symptoms and "angina pectoris" in twins with reference to smoking habits: An epidemiological study with mailed questionnaire. *Arch. Environ. Health* 13:726, 1966 b.
43. CEDERLÖF, R.: Urban factor and prevalence of respiratory symptoms and "angina pectoris": A study on 9,168 twin-pairs with the aid of mailed questionnaires. *Arch. Environ. Health* 13:743, 1966 c.
44. CEDERLÖF, R., JONSSON, E. & LUNDMAN, T.: On the validity of mailed questionnaires in diagnosing "angina pectoris" and "bronchitis". *Arch. Environ. Health* 13:738, 1966 d.
45. CEDERLÖF, R., FRIBERG, L. & JONSSON, E.: Hereditary factors and "angina pectoris". A study on 5,877 twin-pairs with the aid of mailed questionnaires. *Arch. Environ. Health* 14:397, 1967 a.
46. CEDERLÖF, R., EDLUND, M., FRIBERG, L. & JONSSON, E.: Hereditary factors, "spontaneous cough" and "smokers cough": A study on 7,800 twin-pairs with the aid of mailed questionnaires. *Arch. Environ. Health* 14:401, 1967 b.
47. CEDERLÖF, R., FLODERUS, B. & FRIBERG, L.: Cancer in MZ and DZ twins. *Acta Genet. Med. Gemellol.* 19: 69, 1970.
48. CERASI, E. & LUFT, R.: Insulin response to glucose infusion in diabetic and non-diabetic monozygotic twin pairs. Genetic control of insulin response? *Acta Endocrinol.* 55:330, 1967.
49. CHIANG, B.N., PERLMAN, L.V., OSTRANDER, L.D. Jr. & EPSTEIN, F.H.: Relationship of premature systoles to coronary heart disease and sudden death in the Tecumseh epidemiologic study. *Ann. Intern. Med.* 70:1159, 1969.
50. CHIANG, B.N., PERLMAN, L.V., FULTON, M., OSTRANDER, L.D. Jr. & EPSTEIN, F.H.: Predisposing factors in sudden cardiac death in Tecumseh, Michigan. A prospective study. *Circulation* 41:31, 1970.
51. COBB, S., KING, S. & CHEN, E.: Differences between respondents and nonrespondents in a morbidity survey involving clinical examination. *J. Chronic Dis.* 6:95, 1957.
52. CRAMÉR, K., PAULIN, S. & WEIKÖ, L.: Coronary angiographic findings in correlation with age, body weight, blood pressure, serum lipids and smoking habits. *Circulation* 39:888, 1969.
53. DAHLBERG, G.: Twin births and twins from a hereditary point of view. Stockholm, 1926.
54. DAWBER, T.R., MCORE, F.E. & MANN, G.V.: II. Coronary heart disease in the Framingham study. *Am. J. Public Health* 47:4, 1957.
55. DAWBER, T.R. & KANNEL, W.B.: Current status of coronary prevention. Lessons from the Framingham study. *Prev. Med.* 1:290, 1972.
56. DEUTSCHER, S., OSTRANDER, L.D. & EPSTEIN, F.H.: Familial factors in premature coronary heart disease - a preliminary report from the Tecumseh Community Health Study. *Am. J. Epidemiol.* 91:233, 1970.
57. DOAN, A.E., PETERSON, D.R., BLACKMON, J.R. & BRUCE, R.A.: Myocardial ischemia after maximal exercise in healthy men. A method for detecting potential coronary heart disease. *Am. Heart J.* 69:11, 1965.
58. DOLL, R. & HILL, A.B.: "Mortality in relation to smoking: Ten years' observation of British doctors". *Brit. Med. J.* 1:1460, 1964.
59. DOUGLAS, R.
60. DOWNIE, W. and environ
61. DOYLE, J.T. 545, 1970.
62. EDWARDS, J. accuracy of
63. EDWARDS, J.
64. EDMFELDT, mán i jäm
65. EPSTEIN, F.
66. EPSTEIN, F. FRANCIS, T. Ann. Intern.
67. EPSTEIN, F.
68. EPSTEIN, F. Sci. 3:524,
69. FEJFAR, Z.
70. FLETCHER, J.
71. FORSSMAN, aged Swede
72. FOWLER, N.
73. FRANK, C. physical ac
74. FREDRICKS, S.
75. FREDRICKS, S.
76. FRIBERG, L. dizygotic to
77. FRIBERG, L. smoking ha
78. FRIEDMAN, findings, J
79. FRIEDMAN, of the coron
80. FROMENT, et à évolut
81. GEDDA, L. 50 DZ pairs
82. GEDDA, L.
83. GEDDA, L. tosus, and
84. GERTNER, Mass. 1961,
85. GESELL, A.
86. GIKNIS, W. identical to
87. GOFMAN, J. STRISOWER, athero
88. GORDON, J. JAMA Circ
89. GRANT, W. farction in p

59. DOUGLAS, A.H.: Ischemic heart disease in monozygotic twins. *Dis. Chest*, 49:522, 1966.
60. DOWNIE, W.W., BOYLE, J.A., GREIG, W.R., BUCHANAN, W.W. & ALEPA, F.P.: Relative roles of genetic and environmental factors in control of blood pressure in normotensive subjects. *Br. Heart J.* 31:21, 1969.
61. DOYLE, J.T. & KINCH, S.H.: The prognosis of an abnormal electrocardiographic stress test. *Circulation* 41: 545, 1970.
62. EDWARDS, D.A.W., HAMMOND, W.H., HEALY, M.J.R., TANNER, J.M. & WHITEHOUSE, R.H.: Design and accuracy of calipers for measuring subcutaneous tissue thickness. *Br. J. Nutr.* 9:133, 1955.
63. EDWARDS, J.H.: The genetic basis of common disease. *Am. J. Med.* 34:627, 1963.
64. ELMFELDT, D.: Hjärtinfarkt i Göteborg 1968-1970. Morbiditet och mortalitet. Karaktäristika för överlevande män i jämförelse med den manliga befolkningen. Erlanders Boktryckeri, Göteborg, 1974.
65. EPSTEIN, F.H.: Hereditary aspects of coronary heart disease. *Am. Heart J.* 67:445, 1964.
66. EPSTEIN, F.H., OSTRANDER, L.D. Jr., JOHNSON, B.C., PAYNE, M.W., HAYNER, N.S., KELLER, J.B. & FRANCIS, T. Jr.: Epidemiological studies of cardiovascular disease in a total community - Tecumseh, Michigan. *Ann. Intern. Med.* 62:1170, 1965.
67. EPSTEIN, F.H.: Hyperglycemia, A risk factor in coronary heart disease. *Circulation* 36:609, 1967.
68. EPSTEIN, F.H.: Risk factors in coronary heart disease - environmental and hereditary influences. *Isr. J. Med. Sci.* 3:594, 1967.
69. FEJFAR, Z.: "Risk factors" in ischaemic heart disease. *Acta Cardiol. suppl.* 15:7, 1972.
70. FLETCHER, C.M. & HORN, D.: Smoking and health. *WHO Chronicle*, 24:345, 1970.
71. FORSSMAN, O. & LINDEGÅRD, B.: The post coronary patient - a multidisciplinary investigation of middle-aged Swedish males. *J. Psychosom. Res.* 3:89, 1958.
72. FOWLER, N.O.: Clinical diagnosis. *Circulation* 46:1079, 1972.
73. FRANK, C.W., WEINBLATT, E., SHAPIRO, S. & SAGER, R.O.: Myocardial infarction in men. Role of physical activity and smoking in incidence and mortality. *JAMA* 198:1241, 1966.
74. FREDRICKSON, D.S. & LEES, R.S.: A system for phenotyping hyperlipoproteinemia. *Circulation* 31:321, 1965.
75. FREDRICKSON, D.S.: Mutants, hyperlipoproteinaemia, and coronary artery disease. *Br. Med. J.* 2:187, 1971.
76. FRIBERG, L., CEDERLÖF, R., LUNDMAN, T. & OLSSON, H.: Mortality in smoking discordant monozygotic and dizygotic twins. A study on the Swedish Twin Registry. *Arch. Environ. Health* 21:508, 1970.
77. FRIBERG, L., CEDERLÖF, R., LORICH, U., LUNDMAN, T. & deFAIRE, U.: Mortality in twins in relation to smoking habits and alcohol problems. *Arch. Environ. Health* 27:294, 1973.
78. FRIEDMAN, M. & ROSENMAN, R.H.: Association of specific overt behavior pattern with blood and cardiovascular findings. *JAMA* 169:1286, 1959.
79. FRIEDMAN, M. & ROSENMAN, R.H.: Symposium on atherosclerotic heart disease. The prudent management of the coronary-prone individual. *Geriatrics* 27:74, 1972.
80. FROMENT, R., GUINET, P., VIGNON, G. & NOEL, M.: Angor coronarien atheromateux, à debut précoce et à évolution parallèle chez deux jumeaux. *Arch. Mal. Coeur*, 38:260, 1945.
81. GEDDA, L. & POGGI, D.: On the genetic regulation of blood cholesterol. (A study of 50 MZ twin pairs and 50 DZ pairs). *Acta Genet. Med. Gemellol. (Roma)* 9:135, 1960.
82. GEDDA, L.: Twins in history and science. Charles C. Thomas, Springfield, Ill., 1961.
83. GEDDA, L. & BRENCHI, G.: Heredity of biological time and population genetics. Diabetes, lupus erythematosus, and peptic ulcer. *Acta Genet. Med. Gemellol.* 19:493, 1970.
84. GERTLER, M.M. & WHITE, P.D.: Coronary heart disease in young adults: a multidisciplinary study. Cambridge, Mass., 1954. Harvard University Press.
85. GESELL, A.: The method of co-twin control. *Science* 95:446, 1942.
86. GIKNIS, F.L., HOLT, D.E., WHITEMAN, H.W. & SINGH, M.D.: Myocardial infarction in twenty-year-old identical twins. *Am. J. Cardiol.* 16:122, 1965.
87. GOFMAN, J.W., DELALLA, O., GLAZIER, F., FREEMAN, N.K., LINDGREN, F.T., NICHOLS, A.V., STRISOWER, B. & TAMPLIN, A.R.: Serum lipoprotein transport system in health, metabolic disorders, atherosclerosis and coronary heart disease. *Plasma* 2:413, 1954.
88. GORDON, T. & KANNEL, W.B.: Premature mortality from coronary heart disease. The Framingham Study. *JAMA* 215:1617, 1971.
89. GRANT, W.C., WASSERMAN, F., RODENSKY, P.L. & THOMSON, R.V.: The incidence of myocardial infarction in portal cirrhosis. *Ann. Intern. Med.* 51:774, 1959.

90. GRÖNBERG, A., LARSSON, T. & JUNG, J.: Diabetes in Sweden. A clinico-statistical, epidemiological and genetic study of hospital patients and death certificates. *Acta Med. Scand.*, suppl. 477, 1967.
91. GUSTAFSSON, A., ELMFELDT, D., WILHELMSEN, L. & TIBBLIN, G.: Serum lipids and lipoproteins in men after myocardial infarction compared with representative population sample. *Circulation* 46:769, 1972.
92. deHAAS, J.H.: Changing mortality patterns and cardiovascular diseases. Erven Bohn, Haarlem, 1964.
93. HAGERUP, L. & FROM HANSEN, P.: 50-års undersøgelsen i Glostrup 1964-65. *Ugeskr. Læger* 130:1145, 1968.
94. HAMMOND, E.C.: Smoking in relation to the death rates of one million men and women. *Nat. Cancer Inst. Monogr.* 19:127, 1966.
95. HAMMOND, E.C., GARFINKEL, L. & SEIDMAN, H.: Longevity of parents and grandparents in relation to coronary heart disease and associated variables. *Circulation* 43:31, 1971.
96. HARVALD, B. & HAUGE, M.: Heredity in gout and hyperuricemia. *Acta Med. Scand.* 152:247, 1955.
97. HARVALD, B. & HAUGE, M.: Hereditary factors elucidated by twin studies. *Genetics and the Epidemiology of Chronic Disease*, U.S. Department of Health, Education and Welfare, Washington, 1965.
98. HAUGE, M., HARVALD, B., FISCHER, M., GOTLIEB-JENSEN, N., JUEL-NIELSEN, N., RABILD, L., SHAPIRO, R. & VIDEBECK, T.: The Danish twin register. *Acta Genet. Med. (Roma)* 17:315, 1968.
99. HAWK, P.B., OSER, B.L. & SUMMERSON, W.H.: *Practical Physiological Chem.* 13:564, 1954.
100. HELANDER, E.: Om frekvensen av fylleristraf, särskilt hos tvillingar. *Socialmed. tidskr.* 49:555, 1972.
101. HELFANT, R.H., BANKA, V.S., deVILLA, M.A., PINE, R., KABDE, V. & MEISTER, S.G.: Use of bicycle ergometry and sustained handgrip exercise in the diagnosis of presence and extent of coronary heart disease. *Br. Heart J.* 35:1321, 1973.
102. HEYDEN, S., BARTEL, A.G., TABESH, E., CASSEL, J.C., TYROLER, H.A., CORNONI, J.C. & HAMES, C.G.: Angina pectoris and the Rose questionnaire. *Arch. Intern. Med.* 128:961, 1971.
103. HINKLE, L.E. Jr., WHITNEY, L.H., LEHMANN, E.W., DUNN, J., BENJAMIN, B., KING, R., PLAKUN, A. & FLEHINGER, B.: Occupation, education, and coronary heart disease. Risk is influenced more by education and background than by occupational experiences, in the Bell System. *Science* 161:238, 1968.
104. HOFVENDAHL, S. & HELMERS, C.: Ischemisk hjärtsjukdom i olika socio-ekonomiska grupper. *Socialmed. tidskr.* 50:195, 1973.
105. HOLMGREN, A. & STRANDELL, T.: On the use of chest-head leads for recording of electrocardiogram during exercise. *Acta Med. Scand.* 169:57, 1961.
106. HUMERFELT, S.B.: An epidemiological study of high blood pressure, with special reference to the influence of blood pressure and age on cardiac signs and symptoms. *Acta Med. Scand.*, suppl. 407, 1963.
107. International statistical classification of diseases, injuries and causes of death: 1965 revision adapted for indexing of hospital records and morbidity statistics. Stockholm, Civiltryck AB, 1969.
108. JENKINS, C.D., ROSENMAN, R.H. & ZYZANSKI, S.J.: Cigarette smoking: Its relationship to coronary heart disease and related risk factors in the Western Collaborative Group Study. *Circulation* 38:1140, 1968.
109. JENKINS, C.D.: Psychologic and social precursors of coronary disease. First of two parts. *N. Engl. J. Med.* 284:244, 1971.
110. JENKINS, C.D.: Psychologic and social precursors of coronary disease. Second of two parts. *N. Engl. J. Med.* 284:307, 1971.
111. JENSEN, J., BLACKENHORN, D.H., CHIN, H.P., STURGEON, P. & WARE, A.G.: Serum lipids and serum uric acid in human twins. *J. Lipid Res.* 6:193, 1965.
112. JONSELL, S.: A method for the determination of the heart size by teleroentgenography (a heart volume index). *Acta Radiol.* 20:325, 1939.
113. JOSLIN, E.P., ROOT, H., WHITE, P. & MARBLE, A.: *Treatment of diabetes mellitus*. Lea & Febiger, Philadelphia, 1959.
114. KAHLER, O.H. & WEBER, R.: Zur Erbpathologie von Herz- und Kreislaufserkrankungen. Untersuchungen an einer auslesefreien Zwillingserie. II. Coronarsclerose. *Z. Klin. Med.* 137:507, 1940.
115. KAU, L.: Alcoholism in twins. Studies on the etiology and sequelae of abuse of alcohol. Almqvist & Wiksell, Stockholm, 1960.
116. KANNEL, W.B., DAWBER, T.R., KAGAN, A., REVOTSKIE, N. & STOKES, J.: Factors of risk in the development of coronary heart disease - six-year follow-up experience. *Ann. Intern. Med.* 55:33, 1961.
117. KANNEL, W.B.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.

118. KANNEL, W.B.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
119. KANNEL, W.B.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
120. KANNEL, W.B.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
121. KANNEL, W.B.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
122. KANNEL, W.B.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
123. KANNEL, W.B.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
124. KANNEL, W.B.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
125. KASANO, T.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
126. KEMP, J.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
127. KEYS, M.D.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
128. KEYS, M.D.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
129. KEYS, M.D.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
130. KULLER, L.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
131. KULLER, L.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
132. LEADING, J.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
133. LEES, R.S.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
134. LEFSCHE, J.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
135. LEVINE, S.A.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
136. LIKOFF, W.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
137. LIJFHOFF, J.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
138. LUNDMAN, B.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
139. LUNDMAN, B.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
140. MATTING, J.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
141. McDONALD, J.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
142. McKUSICK, V.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
143. MENEMAN, J.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
144. Medical, J.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
145. MYER, J.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.
146. MYER, J.: The Framingham Heart Study: Habits and coronary heart disease. U.S. Public Health Service Publ. No. 1515, 1966.

118. KANNEL, W.B., LeBAUER, E.J., DAWBER, T.R. & McNAMARA, P.M.: Relation of body weight to development of coronary heart disease. The Framingham study. *Circulation* 35:734, 1967.
119. KANNEL, W.B., CASTELLI, W.P. & McNAMARA, P.M.: The coronary profile: 12-year follow-up in the Framingham study. *J. Occup. Med.* 9:611, 1967.
120. KANNEL, W.B., GORDON, T., CASTELLI, W.P. & MARGOLIS, J.R.: Electrocardiographic left ventricular hypertrophy and risk of coronary heart disease. The Framingham study. *Ann. Intern. Med.* 72:813, 1970.
121. KANNEL, W.B., CASTELLI, W.P., GORDON, T. & McNAMARA, P.M.: Serum cholesterol, lipoproteins, and the risk of coronary heart disease. The Framingham study. *Ann. Intern. Med.* 74:1, 1971.
122. KANNEL, W.B., GORDON, T. & SCHWARTZ, M.J.: Systolic versus diastolic blood pressure and risk of coronary heart disease. The Framingham study. *Am. J. Cardiol.* 27:335, 1971.
123. KANNEL, W.B. & FEINLEIB, M.: Natural history of angina pectoris in the Framingham study. Prognosis and survival. *Am. J. Cardiol.* 29:154, 1972.
124. KARVONEN, M.J.: Physical inactivity. In *Preventive Cardiology*, p. 103, 1972.
125. KASANEN, A., KALLIO, V. & FORSGSTRÖM, J.: The significance of psychic and socio-economic stress and other modes of life in the etiology of myocardial infarction. *Ann. Med. Intern. Fenn. suppl.* 143, 1963.
126. KEMP, H.G., ELLIOT, W.C. & GORLIN, R.: The anginal syndrome with normal coronary arteriography. *Trans. Ass. Am. Physicians* 80:59, 1967.
127. KEYS, A. & BLACKBURN, H.: Background of the patient with coronary heart disease. *Progr. Cardio. Dis.* 6:14, 1963.
128. KEYS, A.: Ed. Coronary heart disease in seven countries. *Circulation suppl.* 1, vol. 41, 1970.
129. KEYS, A., ARAVANIS, C., BLACKBURN, H., van BUCHEM, F.S.P., BUZINA, R., DJORDJEVIC, B.S., FIDANZA, F., KARVONEN, M.J., MENOTTI, A., PUDDU, V. & TAYLOR, H.L.: Coronary Heart Disease: Overweight and obesity as risk factors. *Ann. Intern. Med.* 77:15, 1972.
130. KULLER, L., LILIENFELD, A. & FISHER, R.: Epidemiological study of sudden and unexpected deaths due to arteriosclerotic heart disease. *Circulation* 34:1056, 1966.
131. KULLER, L.: Sudden and unexpected non-traumatic deaths in adults: a review of epidemiological and clinical studies. *J. Chronic Dis.* 19:1165, 1966.
132. Leading article. Congenital hyperlipoproteinaemias and ischaemic heart disease. *Lancet* 1:117, 1971.
133. LEES, R.S., CANELLOS, G.P., ROSENBERG, I.H. & HATCH, F.T.: Myocardial infarction in one of a pair of twenty-seven year old identical male twins. *Am. J. Med.* 34:741, 1963.
134. LEPESCHKIN, E. & SUPAWICE, B.: Characteristics of true-positive and false-positive results of electrocardiographic Master two-step exercise tests. *N. Engl. J. Med.* 258:511, 1958.
135. LEVINE, J.B. & ZAK, B.: Automated determination of serum total cholesterol. *Clin. Chim. Acta*, 10:381, 1964.
136. LIKOFF, W., SEGAL, B.L. & KASPARIAN, H.: Paradox of normal selective coronary arteriograms in patients considered to have unmistakable coronary heart disease. *N. Engl. J. Med.* 276:1063, 1967.
137. LILJEFORS, L.: Coronary heart disease in male twins. Hereditary and environmental factors in concordant and discordant pairs. *Acta Med. Scand. suppl.* 511, 1970.
138. LUNDMAN, T.: Smoking in relation to coronary heart disease and lung function in twins. A co-twin control study. *Acta Med. Scand. suppl.* 455, 1966.
139. LUNDMAN, T., LILJEFORS, L., CEDERLÖF, R. & FRIBERG, L.: The validity of the questionnaire diagnosis. *Arch. Environ. Health* 22:597, 1971.
140. MATTINGLY, T.W.: The postexercise electrocardiogram. Its value in the diagnosis and prognosis of coronary arterial disease. *Am. J. Cardiol.* 9:395, 1962.
141. McDONOUGH, J.R., HAMES, C.G., GREENBERG, R.G., GRIFFIN, L.H. Jr. & EDWARDS, A.J. Jr.: Observations on serum cholesterol levels in the twin population of Evans County, Georgia. *Circulation* 25:962, 1962.
142. McKUSICK, V.A. & MURPHY, E.A.: Genetic factors in the etiology of myocardial infarction. In James, T.N. & Keys, J.W., editors: *The etiology of myocardial infarction*, Boston, 1963; Little, Brown and Company.
143. McNEMAR, Q.: *Physiological statistics*, Sec. Ed. Wiley, New York, 1955.
144. Medical Research Council's Committee on the Aetiology of Chronic Bronchitis. Standardized questionnaires on respiratory symptoms. *Br. Med. J.* 2:1665, 1960.
145. MEYER, K.: Serum cholesterol and heredity. A twin study. *Acta Med. Scand.* 172:401, 1962.
146. MIALI, W.E.: Heredity and hypertension. *Practitioner* 207:20, 1971.



147. MOBERG, A.: *Mors subita*, Läkartidningen 66:1011, 1969.
148. MORIYAMA, I.M., KRUEGER, D.E. & STAMLER, J.: Cardiovascular diseases in the United States. Cambridge, Mass., Harvard University Press, 1971.
149. MULCAHY, R., HICKEY, N. & MAURER, B.: Coronary heart disease in women. Study of risk factors in 100 patients less than 60 years of age. *Circulation* 36:577, 1967.
150. MYERS, A.R., EPSTEIN, F.H., DODGE, H.J. & MIKKELSEN, W.M.: The relationship of serum uric acid to risk factors in coronary heart disease. *Am. J. Med.* 45:520, 1968.
151. MYKHED, M.: Alcohol consumption in relation to factors associated with ischemic heart disease. A co-twin control study. *Acta Med. Scand.* suppl. 567, 1974.
152. NAPIER, J.A., METZNER, H. & JOHNSON, B.C.: Limitations of morbidity and mortality data obtained from family histories - a report from the Tecumseh Community Health Study. *Am. J. Public Health* 62:30, 1972.
153. NEWLAND, H.: Antagonism of the anti-thrombotic effect of warfarin by uric acid. *Am. J. Med. Sci.* 256:44, 1968.
154. NIKKILÄ, E.A. & ARO, A.: Family study of serum lipids and lipoproteins in coronary heart-disease. *Lancet* 1:954, 1973.
155. NOBLE, R.P. & CAMPBELL, F.M.: Improved accuracy in automated fluorometric determination of plasma triglycerides. *Clin. Chem.* 16:166, 1970.
156. OLIVER, M.F. & STUART-HARRIS, C.H.: Present position concerning prevention of heart disease. *Br. Med. J.* 2:1203, 1965.
157. OSBORNE, R.H., ADLERSBERG, D., De GEORGE, F.V. & WANG, C.: Serum lipids, heredity and environment. A study of adult twins. *Am. J. Med.* 26:54, 1959.
158. OSBORNE, R.H. & DeGEORGE, F.V.: Genetic basis of morphological variation. An evaluation and application of the twin study method. Harvard University Press. Cambridge, Mass., 1959.
159. OSBORNE, R.H., DeGEORGE, F.V. & MATHERS, J.A.L.: The variability of blood pressure: Basal and casual measurements in adult twins. *Am. Heart J.* 66:176, 1963.
160. OSTRANDER, L.D. Jr., FRANCIS, T. Jr., HAYNER, N.S., KJELSBORG, M.O. & EPSTEIN, F.H.: The relationship of cardiovascular disease to hyperglycemia. *Ann. Intern. Med.* 62:1188, 1965.
161. PARADE, G.W. & LEHMANN, W.: Angina pectoris bei Erbgleichen Zwillingen. *Klin. Wschr.* 17:1036, 1938.
162. PAUL, O. & SCHATZ, M.: On sudden death (Editorial). *Circulation* 43:7, 1971.
163. PENROSE, L.S.: The genetical background of common diseases. *Acta Genet.* 4:257, 1953.
164. PUNSAK, S. & KARVONEN, M.J.: Angina pectoris and ECG abnormalities in relation to prognosis of coronary heart disease in population studies in Finland. In early diagnosis of coronary heart disease. *Adv. Cardiol.* 8: 148, 1973. (Karger, Basel).
165. RAHE, R.H.: Life crisis and health change. From May, Philip R.A., and Wittenborn, J.R., *Psychotropic drug response: Advances in prediction*, 1969, Courtesy of Charles C. Thomas, Springfield, Illin.
166. RAHE, R.H. & LIND, E.: Psychosocial factors and sudden cardiac death: A pilot study. *J. Psychosom. Res.* 15:19, 1971.
167. READER, R. & WYNN, A.: The increasing mortality from coronary heart disease in men, 1950 to 1962. *Med. J. Aust.* 2:740, 1966.
168. REDWOOD, D.R. & EPSTEIN, S.E.: Uses and limitations of stress testing in the evaluation of ischemic heart disease. *Circulation* 46:1115, 1972.
169. RIFKIND, B.M., BOYLE, J.A., GALE, M., GREIG, W. & BUCHANAN, W.W.: Study of serum lipid levels in twins. *Cardiovasc. Res.* 2:148, 1968.
170. ROSE, G.: The diagnosis of ischemic heart pain and intermittent claudication in field surveys. *Bull. WHO* 27:645, 1962.
171. ROSE, G.: Familial patterns in ischaemic heart disease. *Br. J. Prev. Soc. Med.* 18:75, 1964.
172. ROSE, G. & BLACKBURN, H.: Cardiovascular survey methods. WHO Monograph, series, No. 56, 1968.
173. ROSE, G.: Predicting coronary heart disease from minor symptoms and electrocardiographic findings. *Br. J. Prev. Soc. Med.* 25:94, 1971.
174. ROSE, G.: Smoking and cardiovascular disease. *Am. Heart J.* 85:838, 1973.
175. ROSENMAN, R.H., FRIEDMAN, M., STRAUS, R., JENKINS, C.D., ZYZANSKI, S.J. & WURM, M.: Coronary heart disease in the Western Collaborative Group Study. A follow-up experience of 4 1/2 years. *J. Chronic Dis.* 23:173, 1970.
176. Royal College of Physicians, Ltd., 1971.
177. RUSSEK, H.: Coronary heart disease.
178. SALES, S.M.: *Dis.* 23:361, 1971.
179. SELTZER, C.: Coronary heart disease. *JAMA* 225:1000, 1971.
180. SHANOFF, H.: Coronary infarction due to coronary artery disease. *Am. J. Med.* 45:520, 1968.
181. SIDD, J.J., S. N. Engl. J. Med. 281:1000, 1969.
182. SIEVERS, J.: Some clinical observations. *Acta Med. Scand.* 169:95, 1961.
183. SIEVERS, J.: *Acta Med. Scand.* 169:95, 1961.
184. SIMBORG, D.W.
185. SLACK, J. & H. of 121 men and women. *Am. J. Med.* 1:27, 1971.
186. STAMLER, J.: *Am. J. Med.* 1:27, 1971.
187. STAMLER, J.: *Am. J. Med.* 1:27, 1971.
188. STARE, F.J.: *Am. J. Med.* 1:27, 1971.
189. STRANDELL, T.: maximal physical work. *Acta Med. Scand.* 169:95, 1961.
190. SULSER, O.: *Helv. med. Acta* 28:1, 1965.
191. SUNDEY, P.: *Acta Med. Scand.* 169:95, 1961.
192. SYME, S.L.: The occurrence of coronary heart disease. *Adv. Cardiol.* 8: 148, 1973.
193. TAKKUNEN, J.: *Acad. Sci. Fenn. Med. Soc. B* 1971.
194. The Health Council of the Netherlands and Welfare, 1971.
195. The use of twin studies in the study of Gemmology. (D. G. Gemmology, 1971).
196. THENBERG, H.: *Arch. F. Rosen* 1971.
197. THEORELL, T.: variables - a pilot study. *Acta Med. Scand.* 169:95, 1961.
198. THOMAS, C.B.: observations on the use of twin studies. *Acta Med. Scand.* 169:95, 1961.
199. TIBBLIN, G.: *Acta Med. Scand.* 169:95, 1961.
200. TIBBLIN, G.: *Acta Med. Scand.* 169:95, 1961.
201. TIBBLIN, G. & W. *Acta Med. Scand.* 169:95, 1961.
202. TIBBLIN, G.: *Acta Med. Scand.* 169:95, 1961.
203. TIBBLIN, G.: *Acta Med. Scand.* 169:95, 1961.

176. Royal College of Physicians, Smoking and health now. London: Pitman Medical and Scientific Publishing Co. Ltd., 1971.
177. RUSSEK, H.I. & ZOHMAN, B.L.: Relative significance of heredity, diet, and occupational stress in coronary heart disease of young adults. *Am. J. Med. Sci.* 235:266, 1958.
178. SALES, S.M. & HOUSE, J.: Job dissatisfaction as a possible risk factor in coronary heart disease. *J. Chronic Dis.* 23:861, 1971.
179. SELTZER, C.G.: An evaluation of the effect of smoking on coronary heart disease. I. Epidemiological evidence. *JAMA* 203:193, 1968.
180. SHANOFF, H.M., LITTLE, A., MURPHY, E.A. & RYKERT, H.E.: Studies of male survivors of myocardial infarction due to "essential" atherosclerosis. I. Characteristics of the patients. *Can. Med. Assoc. J.* 84:519, 1961.
181. SIDD, J.J., SASAHARA, A.A. & LITTMANN, D.: Coronary-artery disease in identical twins. A family study. *N. Engl. J. Med.* 274:55, 1966.
182. SIEVERS, J., BLOMQVIST, G. & BJÖRCK, G.: Studies on myocardial infarction in Malmö 1935 to 1954: VI. Some clinical data with particular reference to diabetes, menopause and heart rupture. *Acta Med. Scand.* 169:95, 1961.
183. SIEVERS, J.: Myocardial infarction. Clinical features and outcome in three thousand thirty-six cases. *Acta Med. Scand. suppl.* 406, 1963.
184. SIMBORG, D.W.: The status of risk factors and coronary heart disease. *J. Chronic Dis.* 22:515, 1970.
185. SLACK, J. & EVANS, K.A.: The increased risk of death from ischaemic heart disease in first degree relatives of 121 men and 96 women with ischaemic heart disease. *J. Med. Genet.* 3:239, 1966.
186. STAMLER, J. & EPSTEIN, F.H.: Coronary heart disease: Risk factors as guides to preventive action. *Prev. Med.* 1:27, 1972.
187. STAMLER, J.: Epidemiology of coronary heart disease. *Med. Clin. North Am.* 57:1, 1973.
188. STARE, F.J.: Myocardial infarction in patients with portal cirrhosis. *Nutr. Rev.* 19:37, 1961.
189. STRANDELL, T.: Circulatory studies on healthy old men, with special reference to the limitation of the maximal physical working capacity. *Acta Med. Scand. suppl.* 414, 1964.
190. SULSER, O. & KOLLER, F.: Angina pectoris, myocardial infarction and hypercholesteremia in enzygotic twins. *Helv. med. Acta* 28:520, 1961.
191. SUNDEY, P.: Alcoholism and mortality. Universitetsforlaget, Oslo, 1967.
192. SYME, S.L., HYMAN, M.M. & ENTERLINE, P.E.: Some social and cultural factors associated with the occurrence of coronary heart disease. *J. Chronic Dis.* 17:277, 1964.
193. TAKKUNEN, J.: Anthropometric, electrocardiographic and blood pressure studies on adult male twins. *Ann. Acad. Sci. Fenn. (Med)* 107, 1964.
194. The Health Consequences of Smoking. A report to the Surgeon General: 1972. US Dept of Health, Education and Welfare, 1972.
195. The use of twins in epidemiological studies. Report of a WHO meeting of Investigators. *Acta Genet. Med. Gemellol. (Rome)* 15:103, 1966.
196. THEN BERG, H.: Die Erbbiologie des Diabetes Mellitus. Vorläufiges Ergebnis der Zwillingsuntersuchungen. *Arch. F. Essen-u Ges. Biol.* 32:289, 1938.
197. THEORELL, T.: Psychosocial factors in relation to the onset of myocardial infarction and to some metabolic variables - a pilot study. Department of Medicine, Serafimer Hospital, Karolinska Institutet, Stockholm, 1971.
198. THOMAS, C.B. & COHEN, B.H.: Familial occurrence of hypertension and coronary artery disease, with observations concerning obesity and diabetes. *Ann. Intern. Med.* 42:90, 1955.
199. TIBBLIN, G.: A population study of 50-year-old men. An analysis of the non-participation group. *Acta Med. Scand.* 178:453, 1965.
200. TIBBLIN, G.: High blood pressure in men aged 50. A population study of men born in 1913. *Acta Med. Scand. suppl.* 470, 1967.
201. TIBBLIN, G. & WILHELMSEN, L.: Atheroskleros - nikotinsyreterapi (I): Hjärtinfarktens riskfaktorer. *Läkartidningen* 68:1733, 1971.
202. TIBBLIN, G.: Primära och sekundära riskfaktorer. *Läkartidningen* 69:465, 1972.
203. TIBBLIN, G.: Risk factors for developing myocardial infarction and other diseases. The "men born in 1913" study. In: Preventive cardiology (eds. G. Tibblin, A. Keys & L. Werkö), pp 33-42. Almqvist & Wiksell Förlag AB, Stockholm, 1972.

204. TRUETT, J., CORNFELD, J. & KANNEL, W.: A multivariate analysis of the risk of coronary heart disease in Framingham. *J. Chronic Dis.* 20:511, 1967.
205. Twin registries in the study of chronic disease. With particular reference to the relation of smoking to cardiovascular and pulmonary diseases. Report of an International Symposium in San Juan, Puerto Rico, 1-4 Dec. 1969. R. Cederlöf, F.H. Epstein, L. Friberg, Z. Hrubec, E.P. Radford, *Acta Med. Scand.* suppl. 523, 1971.
206. VEDIN, A.J., WILHELMSSON, C., BOLANDER, A.-M. & WERKÖ, L.: Mortality trends in Sweden 1951-1968 with special reference to cardiovascular causes of death. *Acta Med. Scand.* suppl. 515, 1971.
207. VEDIN, A.J., WILHELMSSON, C., ELMFELDT, D., TIBBLIN, G., WILHELMSEN, L. & WERKÖ, L.: Sudden death: Identification of high risk groups. *Am. Heart J.* 86:124, 1973.
208. VERSCHUER, O. von.: Die Zwillingsforschung in Dienste der Inneren Medizin. *Verh. Dtsch. Ges. Inn. Med.* 64:262, 1958.
209. WAHLBERG, F.: Intravenous glucose tolerance in myocardial infarction, angina pectoris and intermittent claudication. *Acta Med. Scand.* suppl. 453, 1966.
210. WARDWELL, W.L., HYMAN, M. & BAHNSON, C.B.: Socio-environmental antecedents to coronary heart disease in 87 white males. *Soc. Sci. Med.* 2:165, 1969.
211. WESTLUND, K. & NICCLAYSEN, R.: Ten-year mortality and morbidity related to serum cholesterol. A follow-up of 3,751 men aged 40-49. *Scand. J. Clin. Lab. Invest.* suppl. 127, 1972.
212. WHITE, P.D.: The genes, the heart and destiny. *N. Engl. J. Med.* 256:965, 1957.
213. WHITE, P.D.: The importance of heredity in coronary heart disease. *Circulation* 22:296, 1960.
214. WIKLAND, B.: Medically unattended fatal cases of ischaemic heart disease in a defined population. Incidence during one year in Stockholm with particular reference to prevalence of certain previously diagnosed disorders, some characteristics of the last attack, and post-mortem findings. *Acta Med. Scand.* suppl. 524, 1971.
215. WILHELMSEN, L., WEDEL, H. & TIBBLIN, G.: Multivariate analysis of risk factors for coronary heart disease. *Circulation* 48:950, 1973.
216. ÅSTRAND, I.: Exercise electrocardiograms recorded twice with an 8-year interval in a group of 204 women and men 48-63 years old. *Acta Med. Scand.* 178:27, 1965.
217. ÅSTRAND, I. and co-workers in the Scandinavian Committee on ECG classification. The "Minnesota code" for ECG classification. Adaptation to CR leads and modification of the code for ECGs recorded during and after exercise. By the Scandinavian Committee on ECG classification. *Acta Med. Scand.* suppl. 481, 1967.
218. ÅSTRAND, I. & LUNDMAN, T.: The exercise electrocardiogram in coronary heart disease. Its prognostic value. *Scand. J. Clin. Lab. Invest.* 22:301, 1968.
219. ÅSTRAND, I.: Exercise ECG, its prognostic significance. *Adv. Cardiol.* 8:142, 1973.

Adapted

#### Q and QS pattern

- (Do not)
- 1.1.1 Q/R amplitude
  - 1.1.2 Q duration
  - 1.1.3 Q duration
  - 1.1.4 Q duration
  - 1.1.5 Q duration
  - 1.1.6 QS pattern
  - 1.1.7 QS pattern
  - 1.2.1 Q/R amplitude leads I, II, III, aVR, aVL, aVF
  - 1.2.2 Q duration
  - 1.2.3 QS pattern
  - 1.2.4 Q duration amplitude
  - 1.2.5 Q duration
  - 1.2.6 Q amplitude
  - 1.2.7 QS pattern
  - 1.2.8 R amplitude CR2 and CR3
  - 1.3.1 Q/R amplitude 0.03 sec
  - 1.3.2 QS pattern
  - 1.3.3 Q duration
  - 1.3.4 Q duration amplitude
  - 1.3.5 Q duration
  - 1.3.6 QS pattern

#### ST depressions

- (Do not)
- 4.1 ST-T depression or 1.0 mV and ST-T depression
  - 4.2 ST-T depression horizontal
  - 4.3 ST-T depression horizontal
  - 4.4 No ST-T depression reaching
  - 4.5 No ST-T depression sloping
  - 4.6 Isolated ST-T depression or 1.0 mV ST segment
  - 4.7 Isolated ST-T depression or 0.5-0.9 mV ST segment

## APPENDIX

The "Minnesota code" for ECG classification.

Adaptation to CR (CH) leads and modification of the code for  
ECGs recorded during and after exercise.

### Q and QS patterns

(Do not code in presence of ventricular conduction defects 6.4 or 7.1)

- 1.1.1 Q/R amplitude ratio 1/3 or more plus Q duration 0.03 sec. or more in any of leads I, II, CR<sub>2</sub>, 3, 4, 5, 6, 7.
- 1.1.2 Q duration 0.04 sec. or more in any of leads I, II, CR<sub>1</sub>, 2, 3, 4, 5, 6, 7.
- 1.1.3 Q duration 0.04 sec. or more plus R amplitude of 3 mm or more in lead aVL.
- 1.1.4 Q duration 0.05 sec. or more in lead III plus any Q wave of at least 1.0 mm amplitude in aVF.
- 1.1.5 Q duration 0.05 sec. or more in lead aVF.
- 1.1.6 QS pattern when R wave is present in adjacent lead to the right on the chest in any of leads CR<sub>2</sub>, 3, 4, 5, 6, 7.
- 1.1.7 QS pattern in all of leads CR<sub>1</sub> through CR<sub>4</sub>, 5, 6 or 7.
- 1.2.1 Q/R amplitude ratio 1/3 or more plus Q duration at least 0.02 sec. and less than 0.03 sec. in any of leads I, II, CR<sub>2</sub>, 3, 4, 5, 6, 7.
- 1.2.2 Q duration at least 0.03 sec. and less than 0.04 sec. in any of leads I, II, CR<sub>2</sub>, 3, 4, 5, 6, 7.
- 1.2.3 QS pattern in lead II.
- 1.2.4 Q duration of at least 0.04 sec. and less than 0.05 sec. in lead III plus any Q wave of at least 1.0 mm amplitude in aVF.
- 1.2.5 Q duration at least 0.04 sec. and less than 0.05 sec. in lead aVF.
- 1.2.6 Q amplitude of 5 mm or more in either of leads III, aVF.
- 1.2.7 QS pattern in all of leads CR<sub>1</sub> through CR<sub>3</sub>.
- 1.2.8 R amplitude decreasing to 2.5 mm or less, and absence of codes 3.2, 7.2 or 7.3 between any of leads CR<sub>2</sub> and CR<sub>3</sub>, CR<sub>3</sub> and CR<sub>4</sub>, CR<sub>4</sub> and CR<sub>5</sub>, CR<sub>5</sub> and CR<sub>6</sub> or CR<sub>6</sub> and CR<sub>7</sub>.
- 1.3.1 Q/R amplitude ratio at least 1/5 and less than 1/3 plus Q duration of at least 0.02 sec. and less than 0.03 sec. in any of leads I, II, CR<sub>2</sub>, 3, 4, 5, 6, 7.
- 1.3.2 QS pattern in absence of code 3.1, in each of leads CR<sub>1</sub> and CR<sub>2</sub>.
- 1.3.3 Q duration of at least 0.03 sec. and less than 0.04 sec. plus R amplitude of 5 mm or more in lead aVL.
- 1.3.4 Q duration of at least 0.03 sec. and less than 0.04 sec. in lead III plus any Q wave of at least 1.0 mm amplitude in lead aVF.
- 1.3.5 Q duration of at least 0.03 sec. and less than 0.04 sec. in lead aVF.
- 1.3.6 QS pattern in each of leads III and aVF.

### ST depressions

(Do not code in presence of ventricular conduction defects 6.4, 7.1, 2.4)

- |  |   |
|--|---|
| <ul style="list-style-type: none"> <li>4.1 ST-J depression of 1.5 mm or more<br/>or 1.0 mm or more<br/>and ST segment straight and slowly ascending, horizontal or downward sloping.</li> <li>4.2 ST-J depression of 1.0-1.4 mm and ST segment straight and slowly ascending,<br/>horizontal or downward sloping.</li> <li>4.3 ST-J depression of 0.5-0.9 mm and ST segment straight and slowly ascending,<br/>horizontal or downward sloping.</li> <li>4.4 No ST-J depression as much as 0.5 mm but ST segment downward sloping and<br/>reaching 0.5 mm or more below P-R baseline.</li> <li>4.5 No ST-J depression as much as 0.5 mm but ST segment horizontal or downward<br/>sloping but reaching less than 0.5 mm below P-R baseline.</li> <li>4.6 Isolated ST-J depression of 1.5 mm or more<br/>or 1.0 mm or more,<br/>ST segment upward sloping.</li> <li>4.7 Isolated ST-J depression of 0.5-1.4 mm or more<br/>or 0.5-0.9 mm or more<br/>ST segment upward sloping.</li> </ul> | <ul style="list-style-type: none"> <li>CR<sub>2</sub>-7,<br/>I, II, aVL, aVF</li> <li>CR<sub>2</sub>-7</li> <li>CR<sub>2</sub>-7<br/>I, II, aVL, aVF</li> <li>CR<sub>2</sub>-7<br/>I, II, aVL, aVF</li> <li>CR<sub>2</sub>-7<br/>I, II, aVL, aVF</li> <li>CR<sub>2</sub>-7<br/>I, II, aVL, aVF</li> <li>CR<sub>2</sub>-7<br/>I, II, aVL, aVF</li> </ul> |
|--|---|

2028553403

2028559404